

Exercise: a “new drug” for elderly patients with chronic heart failure

Roberto Antonicelli^{1*}, Liana Spazzafumo^{2*}, Simonetta Scalvini^{3*}, Fabiola Olivieri^{4,5}, Maria Vittoria Matassini⁶, Gianfranco Parati^{7,8}, Donatella Del Sindaco⁹, Raffaella Gallo¹⁰, and Fabrizia Lattanzio¹¹

¹Department of Cardiology, INRCA-IRCCS National Institute, Ancona, Italy

²Biostatistics Centre, INRCA-IRCCS National Institute, Ancona, Italy

³Department of Cardiology, Maugeri Foundation IRCCS, Lumezzane (BS), Italy

⁴Department of Clinical and Molecular Sciences, DISCLIMO, Università Politecnica delle Marche, Ancona, Italy

⁵Center of Clinical Pathology and Innovative Therapy, INRCA-IRCCS National Institute, Ancona, Italy

⁶Clinic of Cardiology, Università Politecnica delle Marche, Ancona, Italy

⁷Department of Cardiovascular, Neural and Metabolic Sciences, S. Luca Hospital, Italian Auxology Institute, Milano, Italy

⁸Department of Health Sciences, University of Milano-Bicocca, Milan, Italy

⁹Department of Cardiology, INRCA-IRCCS National Institute, Roma, Italy

¹⁰Department of Cardiology, INRCA-IRCCS National Institute, Cosenza, Italy

¹¹Scientific Direction, INRCA-IRCCS National Institute, Ancona, Italy

*These authors equally contributed to this work

Key words: physical exercise, chronic heart failure, quality of life, elderly CHF patients, 6MWT, QoL

Received: 01/15/16; **Accepted:** 01/30/16; **Published:** 03/04/16

Correspondence to: Fabiola Olivieri, PhD; **E-mail:** fabiolaolivieri1@gmail.com

Copyright: Antonicelli et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract: Patients with chronic heart failure (CHF) experience progressive deterioration of functional capacity and quality of life (QoL). This prospective, randomized, controlled trial assesses the effect of exercise training (ET) protocol on functional capacity, rehospitalization, and QoL in CHF patients older than 70 years compared with a control group. A total of 343 elderly patients with stable CHF (age, 76.90±5.67, men, 195, 56.9%) were randomized to ET (TCG, n=170) or usual care (UCG, n=173). The ET protocol involved supervised training sessions for 3 months in the hospital followed by home-telemonitored sessions for 3 months. Assessments, performed at baseline and at 3 and 6 months, included: ECG, resting echocardiography, NT-proBNP, 6-minute walk test (6MWT), Minnesota Living with Heart Failure Questionnaire, and comprehensive geriatric assessment with the InterRAI-HC instrument. As compared to UCG, ET patients at 6 months showed: i) significantly increased 6MWT distance (450±83 vs. 290±97 m, p<0.001); ii) increased ADL scores (5.00±2.49 vs. 6.94±5.66, p=0.037); iii) 40% reduced risk of rehospitalisation (hazard ratio=0.558, 95%CI, 0.326-0.954, p=0.033); and iv) significantly improved perceived QoL (28.6±12.3 vs. 44.5±12.3, p<0.001). In hospital and home-based telemonitored exercise confer significant benefits on the oldest CHF patients, improving functional capacity and subjective QoL and reducing risk of rehospitalisation.

INTRODUCTION

Congestive heart failure (CHF) is a disabling syndrome affecting 1–2% of the adult western population. Its prevalence increases with age, rising to ≥ 10% among individuals aged ≥ 70 years¹ and to 50% among those

aged ≥ 85 years. The proportion of elderly patients with CHF is set to rise as a consequence of three main factors: extended average life expectancy, enhanced response rates to treatments, and longer survival after acute cardiovascular events [2-4].

One of the main consequences of CHF is progressively impaired functional capacity, which in the elderly is exacerbated by a significant amount of comorbidities. Even though emerging evidence shows that exercise in adult CHF patients is associated with improvements in symptoms, exercise tolerance, and quality of life (QoL), which ameliorate clinical outcomes [5], the approach is still far from widespread. However, the data reported by HF-ACTION (Heart Failure: A Controlled Trial Investigating Outcomes of Exercise Training), describes non-significant reductions in the primary endpoint (all-cause mortality or hospitalization) and in key secondary clinical endpoints [6].

A number of clinical trials have confirmed that evidence-based CHF treatments can improve the prognosis of these patients, however very few studies have investigated patients aged > 70 years; in general, HF patients with common comorbidities such as renal failure, liver disease and cognitive impairment were excluded due to their high morbidity and mortality, high rate of hospitalization, and poor QoL [4]. Similarly, few randomized clinical trials, including a small number of patients assessed the safety and effectiveness of cardiac rehabilitation in CHF patients aged more than 75 years [7-10]. One of the earliest trials involving patients older than 75 years, showed an improvement in exercise tolerance and QoL [11].

A recent meta-analysis of seven prospective randomized controlled trials confirmed that exercise training (ET) does not increase mortality/rehospitalisation rates and improves 6 minute-walk test (6MWT) distance in elderly (> 60 years old, range 70-81 years) CHF patients with reduced systolic function and New York Heart Association (NYHA) functional class II or III. The authors conclude that assessment of ET efficacy in old CHF patients requires further investigation using large, rationally designed controlled clinical trials [12].

Recent studies have demonstrated the value of remote telemonitoring in the management of CHF patients [13-18]. However, the effectiveness of telemonitored home-based cardiac rehabilitation exercise in CHF patients aged more than 75 years is still unclear. Interesting results are expected from an ongoing prospective randomized multicentre study of a novel telemedicine approach [19].

This prospective, controlled trial involved stable CHF patients aged more than 70 years randomized to receive a exercise training program in the hospital for 3 months followed by telemonitored home program for the 3 further months, or usual care.

The primary outcome was the effectiveness of CR program in improving functional performance as assessed by the 6MWT; the secondary endpoints were the effectiveness of CR in preventing hospitalization and in improving perceived QoL and multidimensional geriatric assessment.

RESULTS

The flow of the enrolled patients is showed in Figure 1. The 343 patients were randomized to the ET (n=170) and the UC (173) groups. At 3 months (T1), 16 patients dropped out and at 6 months (T2) one more. 13 patients (3.8%) died during the study.

The baseline characteristics of the study population are reported in Table 1.

Men accounted for 56.9% of participants; their mean age was 76.90±5.67 years (43% > 75 years).

As regards the cause of CHF, 49% was due to ischemia, 36% to hypertension, and 15% to valve defects. All patients showed significant physical impairment as assessed by the 6MWT (mean 285±121 m at T0) and had mild to moderate left ventricular dysfunction as measured by a mean left ventricular EF (LVEF) of 48.4% (37% with LVEF < 40%).

There were no differences between the two groups at baseline. The mortality rate during the study was 3.8% (n=13).

Primary endpoint

Functional capacity as measured by 6MWT distance increased significantly in ET at both T1 and T2 ($p < 0.001$) (Table 2). Moreover in the ET group the walked distance compared with baseline was significantly greater at both time points ($p=0.003$).

Secondary endpoints

Over the 6 months of the study, 85 patients (13.1%) required an hospitalisation (25 ET and 60 UC). The effect of ET on the risk of rehospitalisation was significant ($p < 0.001$). The HR, 95% CI, and p values for rehospitalisation for the two groups are reported in Table 3a.

After adjustment for the covariates, ET was found to reduce the risk of all-cause hospitalizations by 44.2% (B coefficient = 0.558, 95%CI, 0.326-0.954, $p=0.033$) (Table 3b, Figure 2a). Classification tree analysis using the variables of the Cox model, i.e. age, gender, BMI, follow-up 6MWT distance, ADL and IADL scores, and

QoL score as dichotomized variables was performed to identify the factors affecting hospitalization (Figure 2b).

BMI < 26.67 involved an increased risk of rehospitalisation in all CHF patients (Figure 2b).




Figure 1. The flow of the 426 consecutive CHF patients enrolled for the study. TCG=training care group; UCG=usual care group.




Figure 2a. Effect of exercise training and usual care on rehospitalization. Cox model adjusted for age, gender, BMI, 6MWT distance, ADL, IADL, and QoL score. TCG=training care group; UCG=usual care group.




Figure 2b. Effect of usual care and exercise training on rehospitalization. Decision-tree analysis developed by CHAID (Chi-squared Automatic Interaction Detector) to assess the risk of rehospitalization including dichotomized variables such as age, gender, BMI, 6MWT distance, ADL, IADL and QoL score as predictors. TCG=training care group; UCG=usual care group. P value adj. = p-value of the chi-square test, adjusted by Bonferroni's correction. DF = degree of freedom.

The InterRAI-HC data are reported in Supplementary Table 1 and show that ADL improved significantly more (p=0.037) in ET than UC participants.

QoL improved significantly in ET compared with UC subjects (p<0.001) (Table 4). Analysis of the respective mean scores showed a comparable perceived QoL at T1 (p=0.987) in both groups, whereas the difference compared with the respective baseline score was sig-

nificant for both groups. The difference between the groups at T2 (ET, 28.6±12.3; UC, 44.5±12.3) was significant (p<0.001).

NT-proBNP was significantly lower (p<0.001) in ET than in UC patients at both time points. The reduction was stronger at T1; however, even though values then rose at T2 in both groups, they remained lower in ET subjects (Table 4).

Table 1. Baseline demographic and clinical characteristics of the groups studied

	TCG (n=170)	UCG (n=173)	Total (n=343)	P value
Age years, mean±SD	76.21±5.21	77.60±6.02	76.90±5.67	0.145
Males , n (%)	103 (60.6%)	92 (53.2%)	195 (56.9%)	0.166
BMI , kg/m ²	26.6±4.4	25.1±5.5	26.8±4.6	0.031
Systolic arterial pressure , mmHg	127.1±15.6	124.9±16.2	126.0±16.0	0.423
Diastolic arterial pressure , mmHg	71.2±9.4	73.6±11.2	72.3±10.3	0.168
6MWT , metres	299±120	270±120	285±121	0.153
LVEF , %	47.9±13.3	49.0±13.4	48.4±13.4	0.166
MLHFQ , score	42.0±14.9	46.8±16.8	44.3±16.0	0.074
NT-proBNP , pg/ml	1236 (2038) [§]	618 (520) [§]	806 (820) [§]	0.110*
Informal support services	170 (100%)	156 (90%)	326 (95%)	<0.001

*Variable was log transformed, § median (interval).

BMI: body mass index, calculated as weight in kilograms divided by the square of the height in metres (kg/m²); LVEF: Left ventricular ejection fraction; 6MWT: Six Minute Walk Test; MLHFQ: Minnesota Living with Heart Failure Questionnaire; NT-proBNP: N-terminal portion of brain-type natriuretic peptide. TCG=training care group; UCG=usual care group. Informal support services: child or child-in-law, other relative, spouse or partner, and friend/neighbour.

Table 2. 6MWT distance in TCG and UCG patients

	TCG	UCG	P value*
6MWT (metres)			
T0	299±120	270±120	<0.001
T1	380.7±120.3	300.6±125.7	
T2	394.1±123.6	301.2±125.8	

*ANOVA for repeated measures. T0: baseline; T1: 3-month follow-up; T2: 6-month follow-up; 6MWT: Six Minute Walk Test; MLHFQ: Minnesota Living with Heart Failure questionnaire; NT-proBNP: N-terminal portion of brain-type natriuretic peptide. TCG=training care group; UCG=usual care group. Continuous data are expressed as mean ± SD and categorical data as number and percentage.

Adherence to and safety of exercise training

Overall, 69% of patients attended at least 70% of hospital rehabilitation sessions; the remaining 31% attended at least 50% of sessions.

As regards telemonitored home ET, 77% of patients performed at least 70% of sessions; the remaining 23% performed at least 50% of sessions.

ET was safe and well tolerated, and sustained arrhythmia, pre-lipothymia and syncope, angina, traumatic events, or falls never occurred during sessions.

DISCUSSION

Whereas the benefits of training have widely been documented in large groups of patients with cardiovascular disease, data on CHF patients, especially on elderly patients, are scarce [20].

Our findings provide fresh evidence for the value of ET in elderly subjects with CHF, by demonstrating that also in this subset of CHF patients the benefits of exercise add to those provided by common medications. In our study, functional capacity improved significantly in ET compared with UC patients, with a significantly greater walked distance at both follow-up assessments in the former group. The 95 m distance increase measured at 6 months is much greater than the previously reported minimal important difference (MID) [21, 22]. The improvement was confirmed by a significant increase in ADL scores. Notably, the ET group experienced a significant reduction in the incidence of all-cause rehospitalisation. The latter finding agrees with the conclusions of recent reviews and meta-analyses that highlighted a significant reduction in the rates of hospitalization of CHF patients performing ET [23]. Finally yet importantly, the improved QoL of our elderly CHF patients is interesting and encouraging, considering that poor physical condition equals social isolation and depression.

Table 3a. Effect of exercise on hospitalization (Cox model)

	TCG (n=150)	UCG (n=163)	P	HR (95%CI)
All-causes hospitalizations	25 (15.2%)	60 (36.8%)	<0.001	2.91 (1.70-4.97)

TCG=training care group; UCG=usual care group.

Table 3b. Effect of exercise on hospitalization, adjusted for clinical covariates

	B	HR	95%CI	P value
UCG vs. TCG	0.583	1.792	1.048-3.065	0.033
Age	-0.017	0.983	0.932-1.036	0.524
Gender	0.129	1.138	0.652-1.985	0.650
ADL	-0.003	0.997	0.934-1.063	0.917
IADL	0.023	1.023	0.965-1.084	0.446
BMI	-0.061	0.941	0.888-0.996	0.037
6MWT	-0.002	0.998	0.996-0.999	0.044
MLHFQ	0.004	1.004	0.989-1.018	0.630

B= beta coefficient. BMI=body mass index; ADL=activities of daily living. IADL=Instrumental activities of daily living. 6MWT=6 Minute Walk Test; MLHFQ=Minnesota Living with the Heart Failure Questionnaire. HR=hazard ratio. TCG=training care group; UCG=usual care group.

Table 4. Change in secondary endpoints (MLHFQ and NT-proBNP) at 3 months and 6 months in the two groups

	TCG	UCG	P value*
MLHFQ			
T0	42.0±14.9	46.8±16.8	<0.001
T1	29.9±9.8	34.7±9.3	
T2	28.6±12.3	44.5±12.3	
NT-proBNP- pg/ml			
T0	1236 (2038) [§]	618 (520) [§]	<0.001
T1	350 (137)	290 (241)	
T2	440 (208)	2143 (1638)	

*ANOVA for repeated measures. T0: baseline; T1: 3-month follow-up; T2: 6-month follow-up; MLHFQ: Minnesota Living with Heart Failure Questionnaire; NT-proBNP: N-terminal portion of brain-type natriuretic peptide. TCG=training care group; UCG=usual care group.

An interesting and innovative aspect of the present study is the use of telemonitoring for the management of cardiologic rehabilitation in elderly CHF patients. We confirmed our previous results on the effects of home telemonitoring in elderly patients with CHF on rate of hospitalization, compliance with treatment and quality of life, by comparison with a group receiving usual care [22, 25].

In the present study, the walked distance was significantly greater, the QL was improved and NT-proBNP was reduced compared with baseline after in hospital ET phase. Notably, these benefits were maintained also after the further telemonitored home phase ET. These results highlights how even the rehabilitation program for patients with CHF followed by remote monitoring has led to significant results. This is of clinical relevance, since the clinical benefit of physical exercise in CHF management is related to the duration of ET. In this context, the telemonitoring of ET at home could ensure adherence to the rehabilitation program over a long period. In our study to increase compliance at home, a nurse tutor call the patients during the home ET sessions, similar to the hospital sessions, and the quality of each patient's ET was asses-

sed by remote telemonitoring of one lead ECG before and after each session. Taking into account the high prevalence of CHF in elderly people, ET telemonitoring at home could be an efficacy tool to ensure compliance with rehabilitation treatment for CHF patients, especially for the oldest old ones.

It was recently reported that the use of telemonitoring in the management of heart failure appears to lead to similar health outcomes as face-to-face or telephone delivery of care [26]. For an elderly subject who is not autonomous in moving from home to the clinic hospital, telemedicine represent a promising prerogative for activities that should be daily.

Conclusion

ET programs associated with drug treatment can attenuate many of the deleterious systemic and tissue-specific effects of CHF, particularly in the elderly. Since CHF is a severe public health problem, particularly in the West, due to progressive ageing of the population, associating physical training and drug treatment can provide important therapeutic effects and thus should be recommended to elderly subjects.

