

Aging and calorie restriction regulate the expression of miR-125a-5p and its target genes *Stat3*, *Casp2* and *Stard13*

Kuldeep Makwana¹, Sonal Arvind Patel¹, Nikkhil Velingkaar¹, Jey Sabith Ebron¹, Girish C. Shukla¹, Roman V. Kondratov¹

¹Center for Gene Regulation in Health and Disease and Department of Biological, Geological, and Environmental Sciences, Cleveland State University, Cleveland, OH 44115, USA

Correspondence to: Roman V. Kondratov; **email:** r.kondratov@csuohio.edu

Keywords: MicroRNA, transcription, calorie restriction, aging, longevity

Received: June 10, 2017 **Accepted:** July 27, 2017 **Published:** July 31, 2017

Copyright: Makwana et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Calorie restriction (CR) is a dietary intervention known to delay aging. In order, to understand molecular mechanisms of CR, we analyzed the expression of 983 MicroRNAs (miRNAs) in the liver of female mice after 2 years of 30% CR using micro-array. 16 miRNAs demonstrated significant changes in their expression upon CR in comparison with age-matched control. mmu-miR-125a-5p (miR-125a-5p) was significantly upregulated upon CR, and in agreement with this, the expression of mRNAs for its three predicted target genes: *Stat3*, *Casp2*, and *Stard13* was significantly downregulated in the liver of CR animals. The expression of precursor miRNA for miR-125a-5p was also upregulated upon CR, which suggests its regulation at the level of transcription. Upon aging miR-125a-5p expression was downregulated while the expression of its target genes was upregulated. Thus, CR prevented age-associated changes in the expression of miR-125a-5p and its targets. We propose that miR-125a-5p dependent downregulation of *Stat3*, *Casp2*, and *Stard13* contributes to the calorie restriction-mediated delay of aging.

INTRODUCTION

Aging, a multifactorial process, is accompanied by the decay of many physiological systems and development of various age-associated pathologies such as cancer, osteoporosis, cardiovascular, and metabolic diseases [1–3]. The human population in developed countries in the world is aged, about 15% of the population in the USA will be older than 65 by the year 2020 [4]. As a consequence of aging, the number of age-associated diseases grows, which leads to increased cost of health care [5]. Therefore, the delay of aging and the increase of health span are important tasks of modern biomedical sciences [6]. Calorie restriction (reduction of calorie intake without malnutrition) is a dietary paradigm, which is known to delay aging and reduce the risk of various age-related pathologies [7,8]. Beneficial effects of CR have been reported in various organisms from

unicellular to mammals [9,10], yet the mechanism of CR is not well deciphered. Increased resistance to stress including oxidative stress, reduced metabolic rate, changes in insulin/IGF/TOR, and sirtuin signaling pathways have been reported and proposed as potential molecular mechanisms of CR [11–14]. Calorie restriction induced changes in gene expression were also proposed as a contributing factor [15,16]. CR can affect gene expression through multiple mechanisms, such as changes in chromatin modifications, mRNA transcription processing, and mRNA translation. Control of the expression of regulatory RNAs such as miRNAs is an important mechanism of gene expression regulation [17–21].

MicroRNAs (miRNAs) are ~22nt long RNA molecules, which incorporate into the RNA-induced silencing complex and regulate gene expression through

sequence-specific interaction with protein-encoding mRNA transcripts. miRNAs induce either mRNA degradation or translational repression [22]. One miRNA may regulate translation of several different transcripts, and the same mRNA can be regulated by several different miRNAs, thus linking many cellular pathways inside the cell [23]. miRNAs are implicated in regulating the process of aging [20,24] expression of some miRNAs is altered in an age-associated manner and can be reversed by calorie restriction [9,19,20,22].

In this study, we investigated the changes in the expression of miRNAs in the liver of mice subjected to a long-term (two years) calorie restriction. miRNA microarray analysis revealed 16 miRNAs with significant changes in their expression pattern upon CR. One of the miRNA, mmu-miR-125a-5p (miR-125a-5p), and its target genes such as *Stat3*, *Casp2*, and *Stard13* demonstrated age-dependent changes, which were ameliorated by CR. Importantly, identified targets have been connected with aging-related signaling pathways. STAT3 is a transcriptional factor that plays a role in the insulin resistance [25], and Casp2 is a proteolytic enzyme, whose activity is upregulated upon aging [26].

RESULTS

CR-induced changes in the liver MicroRNA expression

In order to identify miRNAs affected by CR in the liver of mice, we performed the microarray analysis of miRNAs expression in age-matched female animals fed ad libitum (AL) or 30% CR for two years. miRNAs are known to have daily differential expression pattern [27,28]. Therefore, analysis of miRNA's expression performed at one random time of the day might be misleading. To analyze the possible daily differential expression of miRNAs across the day, we collected liver tissues at six different time points across the day (every four hours) for each group and pooled the samples together for analysis. A microarray containing 983 miRNAs on aµParaflo®Microfluidic Biochip Technology Probe was used for the miRNAs microarray analysis. The hybridization signal intensity for 779 miRNAs was below the threshold level. Therefore they were not considered for further analysis. For the remaining 204 miRNAs, the levels of daily average expression and results of statistical analysis are present-




Figure 1. 30% CR affected the MicroRNA expression in mouse liver. The expression of indicated miRNAs was assayed using (A) microarray hybridization and (B) Taqman PCR assay in the liver of AL (Open bars) and CR mice (Black bars). * - statistically significant difference ($p < 0.05$).

ed in Supplementary Table S1. Among these, CR significantly affected the expressions of 16 miRNAs: 5 miRNAs were down-regulated and 11 were upregulated by CR (see Table 1).

To secondary validate the microarray data, we used TaqMan MicroRNA assay to determine miRNAs expression. We selected two miRNAs: mmu-miR-125a-5p, and mmu-miR-145b (miR-145b) which were found to be significantly affected by CR (Figure 1A). Results of the qPCR analysis are presented in (Figure 1B). The expressions of mature miR-125a-5p and miR-145b were significantly higher in the CR group similar to the microarray data (Figure 1 A & B).

miR-125a-5p is well conserved in mouse and human (see Supplementary Table S3). It is implicated in the development of several pathologies and has a role to play in various tissues upon aging [29–37], We decided to further study the regulation of its expression. We

checked for the expression of precursor transcript of miR-125a-5p. Gene mir-125a is located on the mouse chromosome 17 spanning from 17,830.812 to 17,830,879. This gene encodes for 68 nucleotides long precursor (pre-miR-125a-5p), which further upon Post-transcriptional processing generates miR-125a-5p from its 5' end of the precursor stem. The expression of the precursor was found to be significantly higher in CR samples, shown in (Figure 1C). Thus, both mature MicroRNA and its precursor were upregulated by CR which suggests that the regulation might be controlled at the level of transcription.

CR affected the expression of Mmu-miR-125a-5p target genes

To study the possible physiological consequence of changes in miR-125a-5p expression, we analyzed the effect of CR on its downstream targets. *Stat3* and *Casp2* were previously reported as targets of miR-125a-5p

Table 1. MicroRNAs, which expression was affected by 30% CR in the liver.

Sr.No.	miRNAs	Fold change	p Value	Accession no.
1	mmu-miR-466c-5p	1.6	0.05	MIMAT0004877
2	mmu-miR-145b	1.8	0.02	MIMAT0025105
3	mmu-miR-466m-5p, mmu-miR-669m-5p	1.4	0.04	MIMAT0014882 MIMAT0017346
4	mmu-miR-6240	1.4	0.05	MIMAT0024861
5	mmu-miR-6931-5p	2.1	0.02	MIMAT0027762
6	mmu-miR-125a-5p	1.8	0.003	MIMAT0000135
7	mmu-miR-455-3p	1.6	0.02	MIMAT0003742
8	mmu-miR-139-5p	1.5	0.04	MIMAT0000656
9	mmu-miR-466d-3p	1.6	0.04	MIMAT0004931
10	mmu-miR-5130	2.7	0.04	MIMAT0020641
11	mmu-miR-669h-3p	1.9	0.04	MIMAT0005842
12	mmu-miR-6239	-1.3	0.04	MIMAT0024860
13	mmu-miR-132-3p	-4.7	0.01	MIMAT0000144
14	mmu-miR-6970-3p	-42.9	0.0008	MIMAT0027843
15	mmu-miR-7056-5p	-1.3	0.04	MIMAT0028016
16	mmu-miR-212-3p	-8.1	0.008	MIMAT0000659

[32,38]; using bioinformatics we predicted 4 additional targets: *Nup210*, *Stard13*, *Triap1*, and *Vps4b*. Details of the seeding sequence and target sites on the 3'UTR of identified targets are presented in Supplementary Table S2. Three targets: *Stat3*, *Casp2*, and *Stard13* demonstrated significant downregulation at the mRNA level in CR samples compared with AL (Figure 2). These results were in inverse correlation with the data

on the upregulation of miR-125a-5p in our study. The expression of *Nup210*, *Triap1*, and *Vps4b* genes did not show any statistically significant changes upon CR. Thus, CR induced expression of miRNA caused appropriate changes in the expression of some of its target genes suggesting the potential physiological significance of the regulation of miR-125a-5p expression in response to CR.




Figure 2. CR affected the expression of miR-125a-5p targeted genes in the mouse liver. The expression of indicated mRNAs was assayed in the livers of mice on AL (open bars) or CR (black bars) diets. * - statistically significant difference ($p < 0.05$).




Figure 3. Age dependent changes in the expression of mmu-miR-125a-5p and its target genes. The expressions of (A) mmu-miR-125a-5p, and (B) indicated miR-125a-5p target genes were assayed in the livers of 5 months old (open bars) or 24 month old (black bars) mice. * - statistically significant difference ($p < 0.05$).

CR ameliorated age-induced changes in the expression of miR-125a-5p and its target genes

If miR-125a-5p is relevant to aging, its expression might be affected by age. To directly investigate it we compared the expression of miR-125a-5p and its target genes in the liver of young and old mice. As it is shown in Figure 3A, the expression of mature miR-125a-5p was significantly down-regulated with age. These findings were in agreement with previously reported age-dependent downregulation of mature miR-125a-5p expression in the adipose tissues [18]. In agreement with the reduced expression of miR-125a-5p in the liver of aged mice, the expression of *Stat3*, *Casp2* and *Stard13* was significantly upregulated upon aging (Figure 3B). Thus, aging in mice was accompanied by downregulation of miR-125a-5p expression and by upregulation of the expression of its targets genes such as *Stat3*, *Casp2*, and *Stard13*. The long-term 30% CR ameliorated age-induced changes in miRNA expression of *Stat3*, *Casp2*, and *Stard13*.

To investigate if changes in mRNA level lead to changes at the protein level, we assayed STAT3 protein expression in the liver of young, old, and CR mice. STAT3 expression was upregulated with age in agreement with the upregulation of its mRNA and the downregulation of miR-125a-5p expression (Figure 4A). As it was expected, CR caused a significant reduction in the expression of STAT3 protein in comparison with AL group (Figure 4B). The downregulation of STAT3 expression at the protein level (80% reduction) was stronger than the downregulation

at mRNA level (50% reduction), suggesting that the regulation of the STAT3 expression might occur at both transcript stability and translation levels which are common mechanisms for expression inhibition by miRNAs [39–45].

Stard13 is a direct target of miR-125a-5p

Among three targets affected by CR, *Stat3* and *Casp2* are direct targets of miR-125a-5p [32,38]. Therefore, we decided to test whether the reduction in the expression of *Stard13* upon CR is due to the interaction of mmu-miR-125a-5p with the predicted binding on the 3'UTR of *Stard13*. A region of 3'UTR of *Stard13* containing WT or MUT target site was cloned in the firefly luciferase reporter vector (Figure 5A). In the MUT construct, eight nucleotides in the predicted seed region target site of miR-125a-5p on the 3'UTR of *Stard13* were mutated to disrupt the binding of miR-125a-5p. As shown in Figure 5B, miR-125a-5p mimic reduced the luciferase activity of WT construct by about 60% when compared to cells expressing WT 3'UTR of *Stard13* with co-expression of NC mimic. Whereas, there was no change observed in the luciferase activity for the cells expressing MUT construct either with miR-125a-5p mimic or NC mimic. The repression of firefly luciferase activity as observed in the WT cells co-expressing miR-125a-5p mimic was restored by 50% when the miRNA-mRNA interactions were destabilized in the cells co-expressing MUT construct and miR-125a-5p mimic. This experiment concluded that *Stard13* is a direct target of miR-125a-5p, therefore, explains the downregulation of *Stard13* expression in animals on CR.




Figure 4. CR reversed age induced upregulation of STAT3 protein expression. Representative western blotting and quantification of STAT3 protein expression in the livers of (A) 5 month old (open bars) or 24 months old (black bars) mice. (B) Mice on AL (open bars) or CR (black bars) mice. * - statistically significant difference ($p < 0.05$). Western blot images displayed in the figure were not subjected to any manipulation.




Figure 5. miR125a-5p targets 3'UTR of *Stard13*. (A) luciferase reporter construct containing 228 nucleotide sequence from *Stard13* 3' UTR with either WT or mutant (MUT) miR125a-5p target sites. (B) CHO-K1 cells were cotransfected with the constructs presented in figure 5(A) and with mmu-miR 125a-5p mimic or Negative control (NC) as indicated.

DISCUSSION

miRNAs are the class of regulatory RNAs that are involved in the control of gene expression. miRNAs are implicated in the regulation of many physiological processes [36,38] including aging and several of them have been reported to be affected by age and their expression is changed in response to CR [17], the feeding paradigm known to delay aging. In this study, we have analyzed the expression of miRNAs in the liver of mice who were subject to 2 years 30% CR. Several of the analyzed miRNAs have been previously reported in connection with aging and CR. miR-125a and miR-145 expression declines upon aging in the mouse subcutaneous adipose tissue and CR restore their levels [18]. We have observed similar changes in the expression of these miRNAs in the liver, hence it is reasonable to predict that they can be regulated by aging and CR in different tissues. In agreement with our data, the downregulation of miR-212-3p by CR in the serum of long-lived B6C3F1 mice have been observed [19], however, we did not observe any significant difference in the expression of other 12 miRNAs which were common for both studies [19]. There are many reasons for the discrepancy; including tissue specificity (liver versus serum), time of feeding, time of tissue collection, and mouse strain all as contributing factors. The expres-

sion of miR-93, miR-214, and miR-669c [20-21] in the mouse liver and miR-93 and miR-34a in the rat liver are significantly increased with age [20,46]. In our study, we did not detect any significant effect of CR on miR-214 or miR-669c expression, suggesting that the expression of these two miRNAs is affected by aging but not by CR. The expression of miR-93 and miR-34a were below the detection threshold in our study. The increased expression of mml-miR-451 and mml-miR-144 in the skeletal muscle of old rhesus monkey is reversed by CR [17], but in our study, we did not see any significant effect of CR on the expression of mmu-miR-451, whereas expression of mmu-miR-144 was below the detection threshold. Thus, our and previous study suggest that miRNAs expression is affected by age and diets, but effects are tissue and species specific.

The upregulation of miR-125a-5p expression was also observed at the level of precursor RNA (Figure 1C). EGFR signaling pathway has been reported to be implicated in the regulation of miR-125a-5p expression [35], but to the best of our knowledge, the regulation of miR-125a-5p transcription or post-transcriptional processing has not been studied.

miR-125a-5p is known to be involved in the regulation of cell death [34,47], inflammatory response and lipid

uptake in monocytes/macrophages [31]. miR-125a-5p suppresses tumorigenesis and inhibits metastasis [33,35], which is in agreement with known antitumor effects of CR. Since miRNAs target the expression of multiple genes, we performed bioinformatics analysis, identified several candidates and analyzed the effect of CR on their expression. The expression of *Stat3*, *Stard13*, and *Casp2* was significantly downregulated by CR at mRNA level, and STAT3 was also downregulated at the protein level, which is inversely correlated with the upregulation of miR-125a-5p. In agreement with age-dependent changes in the expression of miR-125a-5p (Figure 3A) the expression of its targets: *Stat3*, *Casp2*, and *Stard13* was increased with age (Figure 3B) and reversed by CR (Figure 2). Therefore, most likely, the regulation of these genes by age and CR occurs in a miRNA-dependent manner. Importantly, *Stat3* and *Casp2* previously reported as targets for miR-125a-5p [32,38].

Effect of aging and CR on STAT3 expression and activity has been previously reported [48,49]. In one study, the increased STAT3 activation (nuclear translocation) in response to growth hormone stimulation has been observed in the liver slices from CR animals compare with slices from AL group, while no significant effect of aging or CR at the total level of STAT3 protein has been detected [48]. In another study, the increase in leptin-stimulated phosphorylation of STAT3 in the hypothalamus was observed in CR group and again no change in the total STAT3 protein level was observed [49]. In our study, we have demonstrated age-dependent upregulation of total STAT3 (Figure 4A) and amelioration of this upregulation upon long-term 30% CR (Figure 3). Together all these results suggest that the effect of diet and age on STAT3 is complicated and might be tissue and treatment specific.

We performed Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway database analysis and identified several pathways relevant to aging, which might be affected by changes in STAT3 level. Chronic activation of STAT3 has been reported to play a part in the development of insulin resistance by increasing the levels of SOCS1 and SOCS3 proteins. SOCS1 and SOCS3, in turn, cause the degradation of insulin receptor substrate IRS-1 and IRS-2, thus, impairing insulin signaling [50–52]. The insulin resistance is one of the manifestations of aging [25] and our data on changes in STAT3 expression with age are in good agreement with age-induced insulin resistance. The prevention of STAT3 induction by CR might contribute to well-known improvement in insulin sensitivity by CR [53,54]. Another potential role of STAT3 in aging revealed by KEGG pathway analysis is the regulation of the RAGE signaling. RAGE is a type I cell surface

receptor for advanced glycation end products (AGE) which might contribute to deregulation of the inflammatory response with age [55,56].

Another target of miR-125a-5p, whose expression was affected by CR, is *Casp2*. CASP2 is a cysteine protease that performs site-specific proteolysis of many intracellular proteins and it is linked to apoptosis [57,58]. The activity of several caspases, including CASP2, has been found to be increased in multiple organs of rats upon aging [26]. It was proposed that the increased caspase activity with age is due to the oxidative stress, and high level of apoptosis in the liver of old MnSOD^{+/−} mice supports this speculation [57–59]. CR inhibits caspase activity which might be due to the reduced oxidative stress [11,60,61] or due to the preventive effect on the age-dependent upregulation of *casp2* as we demonstrated here.

Many age-related pathologies have been associated with the deregulated apoptosis, indeed, failure in the efficient removal of autoreactive or malignant cells lead to the development of auto-immune diseases and cancers. The excessive removal of irreplaceable cells results in organ atrophy and dysfunction, as seen in many aging-associated pathologies like neurodegenerative diseases, kidney diseases, cardiovascular dysfunction and intestinal disorders [61]. Reduced number of hepatocytes and reduction in liver size have been reported upon aging [57,62]. We speculate that via miR-125a-5p mediated manner CR maintains an optimal basal level of CASP2 protein and, thus, maintain basal CASP2 activity to maintain the homeostasis of the liver tissue by decreasing the aging-induced apoptosis.

The third target of miR-125a-5p, whose expression was affected by aging and CR was *Stard13*, also known as *DLC2*. Our study has provided data that suggests *Stard13* to be a direct target of miR-125a-5p (Figure 5A & B). STARD13 is a Rho GTPase-activating protein, which has been suggested to regulate actin cytoskeleton by inactivating Rho GTPases. It was proposed that STARD13 has tumor suppressor functions, which is supported by cell culture-based experiments but not by *in vivo* studies [63,64]. CR has known anti-tumor activity in mammals, hence downregulation of the expression of a potential tumor suppressor gene is counterintuitive, however, *Stard13* might have some other functions, which cannot be ruled out.

There are questions that need to be answered in future studies. What are the molecular mechanisms of the regulation of miRNAs expression by age and by CR? Will the same set of miRNAs be affected in different tissues upon CR? Is it possible to achieve the metabolic effects of CR by modulating the expression of certain

miRNAs? Will the same mechanisms also be functional in primates taking into account that some of identified miRNAs are conserved between rodents and humans? Our results support the importance of miRNA regulation and their pro- and anti-aging activities and warrant further study of identified miRNAs and their targets to understand the mechanism of aging and calorie restriction.

METHODS

Ethics statement

The care and use of mice were carried out in accordance with the guidelines of the Institutional Animal Care and Use Committee (IACUC) of the Cleveland State University. Approval from the Institutional Animal Care and Use Committee (IACUC) of Cleveland State University (Protocol No. 21124-KON-S) was obtained for all the performed animal studies.

Experimental animals

Wild-type mice of C57B6J background were used for the study. Animals were maintained on the 12:12 light: dark cycles with lights turned off and on at 7 am and 7 pm respectively. Animals were fed 18% protein rodent diet (Harlan). The ad libitum (AL) group, had unlimited access to the food all the time. Calorie restriction was implemented gradually. At the age of 3 months, animals were fed 10% calorie restriction diet for the first week i.e. the animals were fed 90% of their total daily food consumption. The second week, animals were maintained on 20% calorie restriction, and after that on 30% calorie restriction diet for two years. The CR group received food as a single meal two hours after the lights were turned off i.e. at ZT14 (Zeitgeber). AL and CR group had unlimited access to water. All the experiments were performed on female mice. At the time of tissue collection, animals were 24 months old.

Cell culture, transfection, and luciferase assay

CHO-K1 cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM: 5% FBS, 2 mM L-glutamine, 1 mM L-proline, 10 mM HEPES) in humidified 5% CO₂ atmosphere at 37°C. For Wild-type (WT) reporter construct, 228bp of 3'UTR of *Stard13* gene spanning putative site for mmu-miR-125a-5p was cloned downstream of firefly luciferase coding region in pMIR-REPORT vector (Ambion, Austin, TX). To generate the mutated construct (MUT), site-directed mutagenesis of the putative target was carried out in the WT-3' UTR construct. Primers used for the cloning and mutagenesis are listed in the Table 2. CHO-K1 cells were plated in 24 well plate one day before the transfection. Cells were co-transfected with 100ng of WT-3'UTR or MUT-3'UTR firefly luciferase reporter construct, 0.5 ng of renilla luciferase reporter plasmid (Promega, Madison, WI) and with either miR-125a-5p mimic (GE lifesciences) or Negative control (NC) mimic (Ambion catalog# 4464058) in duplicates. Cell lysates were collected 48 hours after transfection and assayed for luciferase activity using Dual-Luciferase Reporter Assay System (Promega) and Victor 3 Multilabel Counter 1420 (PerkinElmer). Renilla luciferase activity was used to normalize the firefly luciferase activity.

RNA isolation and mRNA expression analysis

To study the gene expression, liver tissues were collected from six animals, each from AL and CR group. Tissues were collected at six different time points across the day (ZT2, ZT6, ZT10, ZT14, ZT18, and ZT22). Liver tissues collected, were snap frozen on dry ice and stored at -80°C. TriZol reagent was used to isolate the total RNA from the liver tissues according to the manufacturer's protocol (Invitrogen, Carlsbad, CA), and concentration was determined using Nanodrop (Thermo Scientific Nanodrop 2000). RNA expression analysis was performed using real-time RT-PCR with Universal Syber Green mix (BioRad, Hercules, CA) as previously described [65]. Sequences of the primers used for the analysis of expression are listed in the Table 3).

Table 2. Primers used to clone wild type and mutated 3'UTR of Stard13 mRNA.

STARD13 3'UTR	5'-3'
Forward	ATGCACTAGTGCTTTCCCCAGCATGACATC
Reverse	ATGCAAGCTTTAGGCAACGCGGAAGAAATA
Mutant primer	CCAGCATGACATCAAAGAGTCCCTAGAAGAAGGAAAG

Table 3. Primers used for the analysis of mRNA expression for miR-125a-5p target genes.

Gene	Forward (5'-3')	Reverse (5'-3')
Casp2	ACCGTTGAGCTGTGACTATG	GTTCCGTAGCATCTGTGGATAG
Stat3	GTCTGTAGAGCCATACACCAAG	GGTAGAGGTAGACAAGTGGAGA
Nup210	CTCACTCTCATGCCTGTCTATG	GGATTACGGTACTGGATACTG
Vps4b	CAACGGAAGCAAACAACCTCAA	CTCTCTGGCAAGCTGGAATAA
Stard13	CAGGGAAGAAGAAGGGAAAGAA	GGCAACGCGGAAGAAATAAC
Triap1	GGATTAAGGATGGCAGCATTG	GAGCTCCAAAGTTGGTGTTTATG

TaqMan assays for analysis of the mature MicroRNA expression

Taqman assays (ThermoFisher Scientific) were used to validate the results obtained from MicroRNA microarray analysis. Reverse transcription was carried out using reverse transcription kit (ThermoFisher Scientific #4366596) as per the manufacturer's protocol. Total RNA used per reaction for the reverse transcription step was 100ng. Determination of miRNA expression of miR-125a-5p and miR-145b was done by using TaqMan micro RNA assay (ThermoFisher Scientific Catalog # 4427975), and TaqMan Fast Universal PCR master mix 2x (ThermoFisher Scientific Catalog # 4352042) as per the manufacturer's instructions. The detection system used was CFX96 qPCR (BioRad). The reactions for the gene of interest were carried out in triplicates, while normalizing control was run in duplicates. Validation was done using the same RNA samples, which were used for micro array analysis. The comparative delta-Ct method was used to quantify the results. The normalizing control used was snoRNA202 (ThermoFisher Scientific #4427975).

MicroRNA microarray analysis

To assay miRNAs expression profiles, six samples from each AL and CR group were subjected to microarray analysis. Microarray analysis and the data generation was all carried out by LC SCIENCES (Houston, Texas) using μ Paraflo®Microfluidic Biochip Technology. Probe content for all the mature miRNAs was according to the updated version of Sanger miRBase database. All the miRNAs selected, were having a hybridization signal intensity value of 400 or more for an at least one-time point in either of the group.

Immunoblot analysis

Liver tissue samples were immediately frozen on dry ice upon harvesting and stored at -80C before the analysis. Protein extracts were prepared by the lysis of frozen liver tissues in the Cell signaling lysis buffer with the Protease/Phosphatase Inhibitor cocktail (Cell

Signaling Technology, Beverly, MA, USA). Bradford protein assay kit was used to determine the protein concentration using spectrophotometer. Lysates were stored at -80°C. 30ug protein was loaded per well on 4-12% bis-tris gels. Protein was transferred onto the PVDF membrane at 110mA for 70 minutes. The membrane was stained with Ponceau stain to check for the equal loading of proteins. Primary antibody against STAT3 (D1A5) was purchased from Cell Signaling. Staining of the same membranes for beta-actin (SIGMA-ALDRICH #A5441) was used for normalization. All the samples were run on the same gel. Images were obtained with Odyssey FC imaging system (LI-COR, Lincoln, NE), and quantification was performed with Image Studio Lite version 5.2 software.

MicroRNA target site prediction and pathway analysis

Following web resources: Targetscan, miRecords, MicroRNA.org, and mirdb were used to predict targets of individual miRNAs. RNAhybrid - BiBiServ and Targetscan were used to predict the putative target site for miR-125a-5p in the 3'UTR region of the target. KEGG pathway database analysis was done for the targets of miR-125a-5p.

Statistical analysis

For quantitative data, liver tissue samples collected at six different time points across the day for AL and CR groups were run individually. Results are presented as average values plus/minus standard deviation. IBM SPSS Statistics20 and GraphPad Prism Version 5.04software packages were used for statistical analysis. To assay the effect of the diet, unpaired two-tailed T-test was performed. P<0.05 was considered as a statistically significant difference.

Abbreviations

miRNAs: MicroRNA; CR: Calorie Restriction; AL: Adlibitum; miR-125a-5p: mmu-miR-125a-5p; miR-

145b: mmu-miR-145b; Casp2: Caspase2; WT: Wild Type; MUT: Mutant; NC: Negative Control.

AUTHOR CONTRIBUTIONS

K.M., R.V.K., and G.C.S. designed the experiments. Experiments were performed by K.M. K.M. and R.V.K. wrote the manuscript text as well as prepared all the figures. S.P. and N.V., and J.S.E. supplied materials. The manuscript was reviewed by K.M., R.V.K., G.C.S., S.P., J.S.E., and N.V.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

FUNDING

This work is funded by NIH grant 2R01AG039547 and GRHD center at Cleveland State University to RVK. Research in GCS lab is supported by the grants W81XWH-14-1-0508 and W81XWH-14-1-0509 from the department of defense prostate cancer research program.

REFERENCES

1. Feng X, McDonald JM. Disorders of bone remodeling. *Annu Rev Pathol.* 2011; 6:121–45. <https://doi.org/10.1146/annurev-pathol-011110-130203>
2. Rodríguez-Rodero S, Fernández-Morera JL, Menéndez-Torre E, Calvanese V, Fernández AF, Fraga MF. Aging genetics and aging. *Aging Dis.* 2011; 2:186–95.
3. Bonomini F, Rodella LF, Rezzani R. Metabolic syndrome, aging and involvement of oxidative stress. *Aging Dis.* 2015; 6:109–20. <https://doi.org/10.14336/AD.2014.0305>
4. Ortman BJ, Velkoff V a., Hogan H. An Aging Nation: the older population in the United States. *Econ Stat Adm US Dep Commer.* 2014; 1964:1–28.
5. Weatherall D, Greenwood B, Chee HL, Wasi P. Disease Control Priorities in Developing Countries. 2006; 2:119–37.
6. Seals DR, Justice JN, LaRocca TJ. Physiological geroscience: targeting function to increase healthspan and achieve optimal longevity. *J Physiol.* 2016; 594:2001–24. <https://doi.org/10.1113/jphysiol.2014.282665>
7. Colman RJ, Anderson RM, Johnson SC, Kastman EK, Kosmatka KJ, Beasley TM, Allison DB, Cruzen C, Simmons HA, Kemnitz JW, Weindruch R. Caloric restriction delays disease onset and mortality in rhesus monkeys. *Science.* 2009; 325:201–04. <https://doi.org/10.1126/science.1173635>
8. Omodei D, Fontana L. Calorie restriction and prevention of age-associated chronic disease. *FEBS Lett.* 2011; 585:1537–42. <https://doi.org/10.1016/j.febslet.2011.03.015>
9. Lee C, Longo V. Dietary restriction with and without caloric restriction for healthy aging. *F1000 Res.* 2016; 5:F1000 Faculty Rev-117. <https://doi.org/10.12688/f1000research.7136.1>
10. Weindruch R. The retardation of aging by caloric restriction: studies in rodents and primates. *Toxicol Pathol.* 1996; 24:742–45. <https://doi.org/10.1177/019262339602400618>
11. Qiu X, Brown K, Hirschey MD, Verdin E, Chen D. Calorie restriction reduces oxidative stress by SIRT3-mediated SOD2 activation. *Cell Metab.* 2010; 12:662–67. <https://doi.org/10.1016/j.cmet.2010.11.015>
12. Guarente L. Calorie restriction and sirtuins revisited. *Genes Dev.* 2013; 27:2072–85. <https://doi.org/10.1101/gad.227439.113>
13. López-Lluch G, Navas P. Calorie restriction as an intervention in ageing. *J Physiol.* 2016; 594:2043–60. <https://doi.org/10.1113/JP270543>
14. Heilbronn LK, Ravussin E. Calorie restriction and aging: review of the literature and implications for studies in humans. *Am J Clin Nutr.* 2003; 78:361–69.
15. Weindruch R, Kayo T, Lee CK, Prolla TA. Microarray profiling of gene expression in aging and its alteration by caloric restriction in mice. *J Nutr.* 2001; 131:918S–23S.
16. Van Remmen H, Ward WF, Sabia RV, Richardson A. Gene Expression and Protein Degradation. *Handbook of Physiology, Aging.* 2011; 171–234.
17. Mercken EM, Majounie E, Ding J, Guo R, Kim J, Bernier M, Mattison J, Cookson MR, Gorospe M, de Cabo R, Abdelmohsen K. Age-associated miRNA alterations in skeletal muscle from rhesus monkeys reversed by caloric restriction. *Aging (Albany NY).* 2013; 5:692–703. <https://doi.org/10.18632/aging.100598>
18. Mori MA, Raghavan P, Thomou T, Boucher J, Robida-Stubbs S, Macotela Y, Russell SJ, Kirkland JL, Blackwell TK, Kahn CR. Role of microRNA processing in adipose tissue in stress defense and longevity. *Cell Metab.* 2012; 16:336–47. <https://doi.org/10.1016/j.cmet.2012.07.017>
19. Dhahbi JM, Spindler SR, Atamna H, Yamakawa A, Guerrero N, Boffelli D, Mote P, Martin DI. Deep sequencing identifies circulating mouse miRNAs that are functionally implicated in manifestations of aging

- and responsive to calorie restriction. *Aging (Albany NY)*. 2013; 5:130–41.
<https://doi.org/10.18632/aging.100540>
20. Smith-Vikos T, Slack FJ. MicroRNAs and their roles in aging. *J Cell Sci*. 2012; 125:7–17.
<https://doi.org/10.1242/jcs.099200>
 21. Maes OC, An J, Sarojini H, Wang E. Murine microRNAs implicated in liver functions and aging process. *Mech Ageing Dev*. 2008; 129:534–41.
<https://doi.org/10.1016/j.mad.2008.05.004>
 22. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*. 2004; 116:281–97.
[https://doi.org/10.1016/S0092-8674\(04\)00045-5](https://doi.org/10.1016/S0092-8674(04)00045-5)
 23. Carthew RW, Sontheimer EJ. Origins and Mechanisms of miRNAs and siRNAs. *Cell*. 2009; 136:642–55.
<https://doi.org/10.1016/j.cell.2009.01.035>
 24. Kato M, Slack FJ. Ageing and the small, non-coding RNA world. *Ageing Res Rev*. 2013; 12:429–35.
<https://doi.org/10.1016/j.arr.2012.03.012>
 25. Fink RI, Kolterman OG, Griffin J, Olefsky JM. Mechanisms of insulin resistance in aging. *J Clin Invest*. 1983; 71:1523–35.
<https://doi.org/10.1172/JCI110908>
 26. Zhang Y, Chong E, Herman B. Age-associated increases in the activity of multiple caspases in Fisher 344 rat organs. *Exp Gerontol*. 2002; 37:777–89.
[https://doi.org/10.1016/S0531-5565\(02\)00013-X](https://doi.org/10.1016/S0531-5565(02)00013-X)
 27. Figueredo DS, Gitaí DL, Andrade TG. Daily variations in the expression of miR-16 and miR-181a in human leukocytes. *Blood Cells Mol Dis*. 2015; 54:364–68.
<https://doi.org/10.1016/j.bcmd.2015.01.004>
 28. Shende VR, Goldrick MM, Ramani S, Earnest DJ. Expression and rhythmic modulation of circulating microRNAs targeting the clock gene *Bmal1* in mice. *PLoS One*. 2011; 6:e22586.
<https://doi.org/10.1371/journal.pone.0022586>
 29. Lee S, Choi E, Cha MJ, Park AJ, Yoon C, Hwang KC. Impact of miRNAs on cardiovascular aging. *J Geriatr Cardiol*. 2015; 12:569–74. 10.11909/j.issn.1671-5411.2015.05.011
 30. Xu M, Sizova O, Wang L, Su DM. A Fine-Tune Role of Mir-125a-5p on *Foxn1* During Age-Associated Changes in the Thymus. *Aging Dis*. 2017; 8:277–86.
<https://doi.org/10.14336/AD.2016.1109>
 31. Chen T, Huang Z, Wang L, Wang Y, Wu F, Meng S, Wang C. MicroRNA-125a-5p partly regulates the inflammatory response, lipid uptake, and ORP9 expression in oxLDL-stimulated monocyte/macrophages. *Cardiovasc Res*. 2009; 83:131–39.
<https://doi.org/10.1093/cvr/cvp121>
 32. Li D, Kong C, Tsun A, Chen C, Song H, Shi G, Pan W, Dai D, Shen N, Li B. MiR-125a-5p Decreases the Sensitivity of Treg cells Toward IL-6-Mediated Conversion by Inhibiting IL-6R and STAT3 Expression. *Sci Rep*. 2015; 5:14615.
<https://doi.org/10.1038/srep14615>
 33. Hsieh TH, Hsu CY, Tsai CF, Long CY, Chai CY, Hou MF, Lee JN, Wu DC, Wang SC, Tsai EM. miR-125a-5p is a prognostic biomarker that targets HDAC4 to suppress breast tumorigenesis. *Oncotarget*. 2015; 6:494–509.
<https://doi.org/10.18632/oncotarget.2674>
 34. Bhattacharjya S, Nath S, Ghose J, Maiti GP, Biswas N, Bandyopadhyay S, Panda CK, Bhattacharyya NP, Roychoudhury S. miR-125b promotes cell death by targeting spindle assembly checkpoint gene *MAD1* and modulating mitotic progression. *Cell Death Differ*. 2013; 20:430–42.
<https://doi.org/10.1038/cdd.2012.135>
 35. Wang G, Mao W, Zheng S, Ye J. Epidermal growth factor receptor-regulated miR-125a-5p—a metastatic inhibitor of lung cancer. *FEBS J*. 2009; 276:5571–78.
<https://doi.org/10.1111/j.1742-4658.2009.07238.x>
 36. Vickers KC, Remaley AT. MicroRNAs in atherosclerosis and lipoprotein metabolism. *Curr Opin Endocrinol Diabetes Obes*. 2010; 17:150–55.
<https://doi.org/10.1097/MED.0b013e32833727a1>
 37. Barneda-Zahonero B, Parra M. Histone deacetylases and cancer. *Mol Oncol*. 2012; 6:579–89.
<https://doi.org/10.1016/j.molonc.2012.07.003>
 38. Tang L, Shen H, Li X, Li Z, Liu Z, Xu J, Ma S, Zhao X, Bai X, Li M, Wang Q, Ji J. MiR-125a-5p decreases after long non-coding RNA *HOTAIR* knockdown to promote cancer cell apoptosis by releasing caspase 2. *Cell Death Dis*. 2016; 7:e2137.
<https://doi.org/10.1038/cddis.2016.41>
 39. Cimmino A, Calin GA, Fabbri M, Iorio MV, Ferracin M, Shimizu M, Wojcik SE, Aqeilan RI, Zupo S, Dono M, Rassenti L, Alder H, Volinia S, et al. miR-15 and miR-16 induce apoptosis by targeting *BCL2*. *Proc Natl Acad Sci USA*. 2005; 102:13944–49.
<https://doi.org/10.1073/pnas.0506654102>
 40. Poy MN, Eliasson L, Krutzfeldt J, Kuwajima S, Ma X, Macdonald PE, Pfeffer S, Tuschl T, Rajewsky N, Rorsman P, Stoffel M. A pancreatic islet-specific microRNA regulates insulin secretion. *Nature*. 2004; 432:226–30. <https://doi.org/10.1038/nature03076>
 41. Chen X. A microRNA as a translational repressor of *APETALA2* in Arabidopsis flower development. *Science*. 2004; 303:2022–25.
<https://doi.org/10.1126/science.1088060>
 42. Lee RC, Feinbaum RL, Ambros V. The *C. elegans* hetero-

- chronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell*. 1993; 75:843–54. [https://doi.org/10.1016/0092-8674\(93\)90529-Y](https://doi.org/10.1016/0092-8674(93)90529-Y)
43. Bagga S, Bracht J, Hunter S, Massirer K, Holtz J, Eachus R, Pasquinelli AE. Regulation by *let-7* and *lin-4* miRNAs results in target mRNA degradation. *Cell*. 2005; 122:553–63. <https://doi.org/10.1016/j.cell.2005.07.031>
 44. Brennecke J, Hipfner DR, Stark A, Russell RB, Cohen SM. *bantam* encodes a developmentally regulated microRNA that controls cell proliferation and regulates the proapoptotic gene *hid* in *Drosophila*. *Cell*. 2003; 113:25–36. [https://doi.org/10.1016/S0092-8674\(03\)00231-9](https://doi.org/10.1016/S0092-8674(03)00231-9)
 45. Valencia-Sanchez MA, Liu J, Hannon GJ, Parker R. Control of translation and mRNA degradation by miRNAs and siRNAs. *Genes Dev*. 2006; 20:515–24. <https://doi.org/10.1101/gad.1399806>
 46. Li N, Muthusamy S, Liang R, Sarojini H, Wang E. Increased expression of miR-34a and miR-93 in rat liver during aging, and their impact on the expression of *Mgst1* and *Sirt1*. *Mech Ageing Dev*. 2011; 132:75–85. <https://doi.org/10.1016/j.mad.2010.12.004>
 47. Jiang L, Huang Q, Chang J, Wang E, Qiu X. MicroRNA HSA-miR-125a-5p induces apoptosis by activating p53 in lung cancer cells. *Exp Lung Res*. 2011; 37:387–98. <https://doi.org/10.3109/01902148.2010.492068>
 48. Xu X, Sonntag WE. Growth hormone-induced translocation of *stat-3* decreases with age: modulation by calorie restriction. *Am J Physiol*. 1996; 271:34–35.
 49. Wilsey J, Scarpace PJ. Caloric restriction reverses the deficits in leptin receptor protein and leptin signaling capacity associated with diet-induced obesity: role of leptin in the regulation of hypothalamic long-form leptin receptor expression. *J Endocrinol*. 2004; 181:297–306. <https://doi.org/10.1677/joe.0.1810297>
 50. Wunderlich CM, Hövelmeyer N, Wunderlich FT. Mechanisms of chronic JAK-STAT3-SOCS3 signaling in obesity. *JAK-STAT*. 2013; 2:e23878. <https://doi.org/10.4161/jkst.23878>
 51. Kim JH, Kim JE, Liu HY, Cao W, Chen J. Regulation of interleukin-6-induced hepatic insulin resistance by mammalian target of rapamycin through the STAT3-SOCS3 pathway. *J Biol Chem*. 2008; 283:708–15. <https://doi.org/10.1074/jbc.M708568200>
 52. Rui L, Yuan M, Frantz D, Shoelson S, White MF. SOCS-1 and SOCS-3 block insulin signaling by ubiquitin-mediated degradation of IRS1 and IRS2. *J Biol Chem*. 2002; 277:42394–98. <https://doi.org/10.1074/jbc.C200444200>
 53. Prasannarong M, Vichaiwong K, Saengsirisuwan V. Calorie restriction prevents the development of insulin resistance and impaired insulin signaling in skeletal muscle of ovariectomized rats. *Biochim Biophys Acta*. 2012; 1822:1051–61. <https://doi.org/10.1016/j.bbadis.2012.02.018>
 54. Larson-Meyer DE, Heilbronn LK, Redman LM, Newcomer BR, Frisard MI, Anton S, Smith SR, Alfonso A, Ravussin E. Effect of calorie restriction with or without exercise on insulin sensitivity, beta-cell function, fat cell size, and ectopic lipid in overweight subjects. *Diabetes Care*. 2006; 29:1337–44. <https://doi.org/10.2337/dc05-2565>
 55. Yamagishi S, Matsui T. Role of receptor for advanced glycation end products (RAGE) in liver disease. *Eur J Med Res*. 2015; 20:15. <https://doi.org/10.1186/s40001-015-0090-z>
 56. Riehl A, Németh J, Angel P, Hess J. The receptor RAGE: bridging inflammation and cancer. *Cell Commun Signal*. 2009; 7:12. <https://doi.org/10.1186/1478-811X-7-12>
 57. Lopez-Cruzan M, Herman B. Loss of caspase-2 accelerates age-dependent alterations in mitochondrial production of reactive oxygen species. *Biogerontology*. 2013; 14:121–30. <https://doi.org/10.1007/s10522-013-9415-x>
 58. Zhang JH, Zhang Y, Herman B. Caspases, apoptosis and aging. *Ageing Res Rev*. 2003; 2:357–66. [https://doi.org/10.1016/S1568-1637\(03\)00026-6](https://doi.org/10.1016/S1568-1637(03)00026-6)
 59. Kokoszka JE, Coskun P, Esposito LA, Wallace DC. Increased mitochondrial oxidative stress in the *Sod2* (+/-) mouse results in the age-related decline of mitochondrial function culminating in increased apoptosis. *Proc Natl Acad Sci USA*. 2001; 98:2278–83. <https://doi.org/10.1073/pnas.051627098>
 60. Lee JH, Jung KJ, Kim JW, Kim HJ, Yu BP, Chung HY. Suppression of apoptosis by calorie restriction in aged kidney. *Exp Gerontol*. 2004; 39:1361–68. <https://doi.org/10.1016/j.exger.2004.06.015>
 61. Phelouzat MA, Quadri RA, Proust JJ. Apoptosis and Aging. *M S-Medecine Sciences*. 5th ed. 1995; 894–900.
 62. Thomas RP, Guigneaux M, Wood T, Evers BM. Age-associated changes in gene expression patterns in the liver. *J Gastrointest Surg*. 2002; 6:445–53. [https://doi.org/10.1016/S1091-255X\(01\)00010-5](https://doi.org/10.1016/S1091-255X(01)00010-5)
 63. Yau TO, Leung TH, Lam S, Cheung OF, Tung EK, Khong PL, Lam A, Chung S, Ng IO. Deleted in liver cancer 2 (DLC2) was dispensable for development and its deficiency did not aggravate hepatocarcinogenesis. *PLoS One*. 2009; 4:e6566.

<https://doi.org/10.1371/journal.pone.0006566>

64. Tang F, Zhang R, He Y, Zou M, Guo L, Xi T. MicroRNA-125b induces metastasis by targeting STARD13 in MCF-7 and MDA-MB-231 breast cancer cells. *PLoS One*. 2012; 7:e35435.

<https://doi.org/10.1371/journal.pone.0035435>

65. Patel SA, Velingkaar N, Makwana K, Chaudhari A, Kondratov R. Calorie restriction regulates circadian clock gene expression through BMAL1 dependent and independent mechanisms. *Sci Rep*. 2016; 6:25970.

<https://doi.org/10.1038/srep25970>

SUPPLEMENTARY MATERIAL

Supplementary Table S1. The effect of 30% CR on miRNA expression in the liver.

	Group 1		Group 2		
	Cr		Ad		
Reporter Name	Mean	StDev	Mean	StDev	T-test(p-Value)
mmu-miR-3960	7,214	3,720	5,893	1,870	0.345729
mmu-miR-7028-5p	10,850	5,504	11,422	12,795	0.707084
mmu-miR-133b-5p	29,166	4,790	27,453	17,687	0.996587
mmu-let-7k	347	174	359	360	0.770206
mmu-miR-3962	317	180	368	322	0.465436
mmu-miR-3620-5p	2,201	1,202	2,010	944	0.806528
mmu-miR-669c-5p	1,567	488	1,633	403	0.863513
mmu-miR-2861	2,335	1,637	1,198	426	0.039967
mmu-miR-669f-5p	939	270	934	236	0.810106
mmu-miR-6238	615	747	2,960	4,161	0.048597
mmu-miR-341-5p	56,348	8,592	47,700	12,934	0.012713
mmu-miR-101b-3p	1,044	446	1,213	426	0.326458
mmu-miR-100-5p	319	96	438	247	0.091844
mmu-miR-143-3p	1,202	628	911	121	0.131428
mmu-miR-27a-3p	1,034	309	1,105	321	0.494768
mmu-miR-15a-5p	968	470	1,402	576	0.036146
mmu-miR-669o-5p	396	150	354	105	0.454633
mmu-miR-195a-5p	496	315	490	155	0.984438
mmu-miR-6937-5p	4,972	2,040	4,313	1,551	0.506479
mmu-miR-669l-5p	419	149	380	111	0.502784
mmu-miR-669n	1,609	472	1,693	423	0.773574
mmu-miR-466g	1,736	510	1,625	366	0.559514
mmu-miR-7047-5p	3,664	1,589	3,539	1,266	0.978717
mmu-miR-466c-5p	599	206	365	139	0.006519
mmu-miR-24-3p	3,017	1,225	2,871	309	0.641511
mmu-miR-194-5p	6,894	1,425	7,668	1,980	0.318193
mmu-miR-149-3p	3,665	1,945	3,240	1,146	0.656297
mmu-miR-25-3p	512	235	469	85	0.506317
mmu-miR-3099-3p	564	144	561	125	0.918921
mmu-let-7j	511	131	557	251	0.504248
mmu-miR-466j	409	205	574	291	0.235113
mmu-miR-101c	274	173	303	292	0.601243
mmu-miR-2137	1,320	732	915	382	0.127494
mmu-miR-148a-3p	2,704	672	3,054	912	0.241186
mmu-miR-5100	1,232	940	700	95	0.077305
mmu-miR-762	5,644	1,968	5,078	2,189	0.629904
mmu-let-7e-5p	273	151	234	252	0.811858
mmu-miR-30e-5p	1,994	950	2,014	604	0.860057
mmu-miR-466h-5p	265	185	504	262	0.031873
mmu-miR-3473b	875	534	465	216	0.037754
mmu-miR-669k-5p	387	154	383	81	0.86831
mmu-miR-466h-3p	3,202	708	3,282	558	0.878888
mmu-miR-3095-3p	877	192	826	189	0.531237
mmu-miR-466i-5p	3,884	832	4,156	1,033	0.674214
mmu-miR-30a-5p	3,669	1,104	3,600	731	0.907727
mmu-miR-145b	893	219	473	288	0.001227
mmu-miR-466m-5p	926	269	632	134	0.003522

mmu-miR-29b-3p	894	666	1,157	637	0.264488
mmu-miR-151-5p	334	135	334	124	0.986799
mmu-miR-6970-5p	512	314	399	154	0.35769
mmu-miR-27b-3p	1,737	431	1,888	355	0.45367
mmu-miR-1934-3p	399	266	277	154	0.244947
mmu-miR-221-3p	786	251	735	162	0.519058
mmu-miR-3072-5p	385	123	455	167	0.247301
mmu-miR-669c-3p	2,285	623	2,092	361	0.432303
mmu-miR-3077-5p	392	327	221	104	0.125798
mmu-miR-669f-3p	1,807	479	1,663	241	0.420768
mmu-miR-6240	502	107	345	128	0.00734
mmu-miR-3970	68	41	223	257	0.034218
mmu-miR-5126	8,401	3,751	6,718	1,756	0.206182
mmu-miR-7683-3p	136	135	318	369	0.096302
mmu-miR-3535	395	204	254	123	0.048215
mmu-let-7g-5p	3,091	535	2,883	715	0.330805
mmu-miR-6239	14,009	3,189	18,268	3,055	0.00601
mmu-miR-6931-5p	801	377	366	108	0.001608
mmu-miR-378c	329	127	355	83	0.772099
mmu-miR-6385	425	140	443	89	0.783923
mmu-miR-485-3p	529	127	572	231	0.861091
mmu-miR-101a-3p	1,160	558	1,346	555	0.357532
mmu-miR-378a-3p	446	149	514	129	0.396116
mmu-miR-6366	277	138	266	75	0.865699
mmu-miR-26b-5p	2,069	582	1,718	405	0.111709
mmu-miR-3473f	488	363	199	76	0.021179
mmu-let-7f-5p	3,119	892	2,538	680	0.092733
mmu-miR-7045-5p	1,231	502	1,132	268	0.638869
mmu-miR-23b-3p	2,288	634	2,160	503	0.508675
mmu-miR-705	870	350	736	199	0.344456
mmu-miR-3082-5p	1,533	430	1,541	380	0.937836
mmu-miR-1892	557	337	521	256	0.839428
mmu-miR-3963	1,215	695	1,138	366	0.861965
mmu-miR-6908-5p	272	110	372	124	0.081446
mmu-miR-22-3p	7,143	1,381	7,467	1,743	0.688749
mmu-miR-5099	2,857	903	2,234	497	0.056274
mmu-miR-20b-5p	193	101	289	171	0.072652
mmu-miR-466q	1,750	494	1,560	372	0.346314
mmu-miR-7011-5p	340	117	435	96	0.039827
mmu-miR-6965-3p	689	371	2,111	2,373	0.037218
mmu-miR-423-5p	319	59	353	188	0.492977
mmu-miR-107-3p	1,048	372	1,331	287	0.08274
mmu-miR-329-3p	34	20	144	316	0.210463
mmu-miR-7221-3p	1,527	853	1,031	348	0.106025
mmu-miR-30c-5p	5,378	1,120	5,293	1,404	0.797422
mmu-miR-466f	526	197	537	212	0.779986
mmu-miR-145a-5p	1,726	554	1,417	209	0.087719
mmu-let-7d-5p	2,497	735	2,091	837	0.265377
mmu-miR-20a-5p	819	292	805	182	0.88494
mmu-miR-17-5p	558	209	616	187	0.481723
mmu-miR-363-5p	132	84	409	327	0.008187
mmu-miR-21a-5p	5,170	2,064	4,156	1,694	0.148317
mmu-miR-466f-3p	3,338	638	3,353	456	0.988181
mmu-miR-5130	332	215	120	48	0.004882
mmu-miR-7082-5p	2,063	640	2,862	1,106	0.0436

mmu-miR-378b	350	147	371	90	0.871018
mmu-miR-466i-3p	2,524	582	2,427	388	0.626978
mmu-miR-30d-5p	2,078	619	1,832	851	0.539806
mmu-miR-466f-5p	483	153	373	129	0.024391
mmu-miR-5119	277	189	218	118	0.451497
mmu-miR-6929-3p	326	72	368	110	0.387556
mmu-miR-6981-5p	941	263	1,165	229	0.044817
mmu-miR-130a-3p	967	286	1,011	223	0.777342
mmu-miR-93-5p	434	182	458	65	0.70582
mmu-let-7i-5p	845	236	864	270	0.89157
mmu-miR-7658-5p	248	191	118	40	0.039922
mmu-miR-1187	1,765	391	2,275	725	0.084365
mmu-miR-5107-5p	429	174	648	210	0.008225
mmu-miR-346-3p	626	302	468	183	0.21009
mmu-miR-122-3p	2,205	817	2,301	1,003	0.780025
mmu-miR-669e-5p	154	124	287	151	0.055
mmu-miR-342-3p	160	82	231	153	0.139103
mmu-miR-29c-3p	2,844	1,239	3,114	1,290	0.501129
mmu-miR-3473e	702	435	333	144	0.019497
mmu-miR-152-3p	537	190	553	191	0.830174
mmu-miR-5113	512	135	468	134	0.473942
mmu-miR-22-5p	479	165	506	138	0.764314
mmu-miR-214-3p	546	374	536	92	0.948595
mmu-miR-103-3p	1,064	398	1,298	295	0.180138
mmu-let-7a-5p	2,818	843	2,284	920	0.193003
mmu-miR-802-5p	412	256	386	250	0.935593
mmu-miR-6944-5p	660	359	512	201	0.328479
mmu-miR-181a-5p	419	533	207	24	0.200903
mmu-miR-6896-3p	621	147	679	183	0.487437
mmu-miR-8101	315	160	306	109	0.956405
mmu-miR-494-3p	306	108	607	455	0.020016
mmu-miR-690	12,388	2,556	12,478	1,637	0.865777
mmu-miR-466m-3p	1,589	411	1,530	227	0.683266
mmu-miR-146a-5p	571	222	642	231	0.454848
mmu-miR-6922-5p	492	201	502	116	0.907653
mmu-miR-669a-3p	1,975	540	1,877	276	0.645677
mmu-miR-1895	875	202	1,151	258	0.012891
mmu-miR-30b-5p	3,305	908	3,238	856	0.850504
mmu-miR-672-5p	245	137	441	238	0.041637
mmu-miR-92b-3p	420	175	315	64	0.086432
mmu-miR-203-3p	453	118	519	178	0.39029
mmu-miR-8093	54	60	138	285	0.287563
mmu-miR-3097-5p	377	157	397	231	0.854942
mmu-miR-99a-5p	863	298	887	402	0.779772
mmu-miR-31-5p	749	226	759	158	0.958095
mmu-miR-669e-3p	1,689	440	1,579	195	0.529697
mmu-miR-122-5p	10,790	972	11,909	2,243	0.237239
mmu-miR-1224-5p	419	135	432	91	0.707676
mmu-miR-23a-3p	2,104	569	1,890	487	0.323776
mmu-miR-5112	4,114	2,707	6,888	5,212	0.071404
mmu-miR-7661-3p	280	96	388	91	0.018606
mmu-let-7c-5p	2,337	706	1,949	839	0.276307
mmu-miR-451a	4,801	844	4,358	1,396	0.431628
mmu-miR-32-3p	785	266	719	202	0.528477
mmu-miR-669a-3-3p	739	278	579	153	0.131556

mmu-miR-7239-3p	3,747	4,414	8,162	10,140	0.133449
mmu-miR-15b-5p	412	152	321	82	0.092767
mmu-miR-7027-3p	283	53	296	106	0.764451
mmu-miR-19b-3p	492	262	562	197	0.594425
mmu-miR-26a-5p	5,655	751	5,397	1,012	0.302324
mmu-miR-7081-5p	3,523	2,029	5,329	5,554	0.203969
mmu-miR-125a-5p	840	158	446	194	4.533E-5
mmu-miR-467a-3p	1,210	378	1,079	286	0.412676
mmu-miR-5121	518	205	392	82	0.044916
mmu-miR-7235-3p	594	93	649	180	0.463036
mmu-miR-467f	2,627	602	2,567	465	0.768961
mmu-miR-714	413	245	250	80	0.062626
mmu-let-7b-5p	1,444	466	1,290	624	0.585893
mmu-miR-467e-3p	740	248	593	164	0.143019
mmu-miR-467c-3p	880	293	736	194	0.220199
mmu-miR-574-5p	2,597	552	3,040	938	0.295107
mmu-miR-483-5p	1,085	329	1,260	261	0.182766
mmu-miR-212-3p	32	12	259	171	0.000167
mmu-miR-6348	1,258	426	1,403	400	0.542885
mmu-miR-467g	701	242	512	172	0.053295
mmu-miR-574-3p	1,659	415	1,475	342	0.311189
mmu-miR-126a-3p	4,832	1,128	4,781	914	0.920253
mmu-miR-7016-5p	269	63	363	209	0.112953
mmu-miR-467b-3p	2,685	618	2,764	444	0.805065
mmu-miR-199a-5p	342	295	301	45	0.670773
mmu-miR-468-3p	428	145	357	159	0.113217
mmu-miR-3569-5p	262	75	337	93	0.029548
mmu-miR-467d-3p	2,044	547	2,004	398	0.865427
mmu-miR-192-5p	6,114	1,229	6,908	1,594	0.294265
mmu-miR-7116-5p	285	368	396	290	0.335222
mmu-miR-199a-3p	605	491	625	168	0.888147
mmu-miR-700-3p	259	86	329	184	0.192203
mmu-miR-191-5p	1,121	397	1,162	270	0.812239
mmu-miR-200b-3p	181	161	113	45	0.156052
mmu-miR-1191b-5p	106	131	129	275	0.705775
mmu-miR-328-5p	274	91	319	119	0.214503
mmu-miR-455-3p	371	108	231	71	0.00106
mmu-miR-200a-3p	222	205	145	55	0.210269
mmu-miR-16-5p	4,536	997	5,326	875	0.069498
mmu-miR-709	13,326	2,032	16,085	2,474	0.012694
mmu-miR-8090	53	75	138	334	0.35321
mmu-miR-92a-3p	800	306	670	132	0.170368
mmu-miR-7056-5p	347	99	484	112	0.002219
mmu-miR-139-5p	261	90	166	43	0.004586
mmu-miR-466d-3p	260	111	153	39	0.008479
mmu-miR-6970-3p	47	15	2,013	1,567	0.000219
mmu-miR-132-3p	73	38	345	371	0.014901

