

***In vitro* P38MAPK inhibition in aged astrocytes decreases reactive astrocytes, inflammation and increases nutritive capacity after oxygen-glucose deprivation**

Miren Revuelta^{1,2,#}, Amaia Elicegui^{1,3}, Till Scheuer¹, Stefanie Endesfelder¹, Christoph Bühner¹, Leire Moreno-Cugnon^{2,*}, Ander Matheu^{2,4,5}, Thomas Schmitz¹

¹Department for Neonatology, Charité University Medical Center, Berlin 13353, Germany

²Cellular Oncology Group, Biodonostia Health Research Institute, Paseo Doctor Begiristain, San Sebastian 20014, Spain

³Neurovascular Research Laboratory, Vall d'Hebron Institute of Research, Barcelona 08035, Spain

⁴IKERBASQUE, Basque Foundation for Science, Bilbao 48013, Spain

⁵CIBERfes, Madrid 28029, Spain

*Current address: Nerve Disorders Laboratory, CIC bioGUNE, Derio, Vizcaya, Spain

#Current address: Department of Physiology, Faculty of Pharmacy, University of the Basque Country UPV/EHU, Vitoria, Spain

Correspondence to: Thomas Schmitz, Miren Revuelta; **email:** thomas.schmitz@charite.de, miren.revuelta@ehu.eus

Keywords: astrocytes, OGD, ageing, p38MAPK

Received: July 22, 2020

Accepted: December 18, 2020

Published: February 9, 2021

Copyright: © 2021 Revuelta et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/3.0/) (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Proper astroglial functioning is essential for the development and survival of neurons and oligodendroglia under physiologic and pathological circumstances. Indeed, malfunctioning of astrocytes represents an important factor contributing to brain injury. However, the molecular pathways of this astroglial dysfunction are poorly defined. In this work we show that aging itself can drastically perturb astrocyte viability with an increase of inflammation, cell death and astrogliosis. Moreover, we demonstrate that oxygen glucose deprivation (OGD) has a higher impact on nutritive loss in aged astrocytes compared to young ones, whereas aged astrocytes have a higher activity of the anti-oxidant systems. P38MAPK signaling has been identified to be upregulated in neurons, astrocytes and microglia after ischemic stroke. By using a pharmacological p38 α specific inhibitor (PH-797804), we show that p38MAPK pathway has an important role in aged astrocytes for inflammatory and oxidative stress responses with the subsequent cell death that occurs after OGD.

INTRODUCTION

Ischemic stroke, the most common type of stroke, represents the second main cause of death worldwide leading to 5.7 million of deaths per year in adulthood [1, 2]. Between all types of stroke, brain ischemic stroke represent 87% of all cases that cause neurological deficits such as motor impairment and inability to read or even aphasia [3].

During the last decades, ischemic stroke experimental models have been performed in young animals, but aging itself plays a critical role in the response of the brain to stroke [4]. Indeed, brain aging is accompanied by many structural and physiological alterations that usually involve cognitive decline [5]. Although most of the molecular mechanisms of brain injury are similar in neonatal and aged animal models [6, 7], certain characteristics are different, e.g., cell apoptosis

occurs a week after the injury in the immature brain while this period takes place only for a few hours after the damage in the mature brain [8]. For *in vitro* studies, the commonly used model to mimic cerebral ischemia is the oxygen glucose deprivation model (OGD) [9, 10].

Although neurons are primarily susceptible to injury, impairment of supporting glial cells such as astrocytes may contribute to secondary injury in neurons [11]. Proper astroglial functioning is essential for the development and survival of neurons and oligodendroglia under pathological circumstances [12]. Supportive properties of astrocytes are reflected in production of growth factors such as PDGF and IGF [13, 14], clearance of radicals via superoxide dismutase activity [15], anti-oxidant defense with glutathione synthesis [16], removal of glutamate from the synaptic cleft [17], among others. Hence, malfunctioning of astrocytes represents an important factor for recovery and repair of the brain after injury [18, 19]. The importance of age for differences in glial cellular responses after brain injury were also documented in microglia [20].

p38 mitogen-activated protein kinase (p38MAPK) pathway is a signaling pathway that can be activated in neurodegenerative diseases [21, 22]. p38MAPK controls key processes of mammalian cell homeostasis such as self-renewal, differentiation, proliferation and death [23]. Activation of p38MAPK signaling has been identified in neurons, astrocytes and microglia after ischemic stroke [24]. Moreover, p38MAPK inhibitors seem to be effective reducing infarct volume after stroke [21, 25, 26], but the cell type being involved in p38MAPK dependent injury in the brain after stroke remains to be elucidated, as well as the effect of p38MAPK in aged cells after stroke.

Therefore, our aim was to investigate the effect of OGD in young and aged primary rat astrocyte cultures and to analyze the expression and effect of p38MAPK in these cultures. For this, we cultured young and aged astrocytes with PH-797804, MAPK14 (*p38 α*) inhibitor, the most abundant isoform of p38MAPK in the brain, to define changes in supporting and protective properties of astrocytes that can be critical for survival of brain cells.

MATERIALS AND METHODS

Animals

All animal experiments were performed in accordance to German and Spanish animal welfare law with the permission of the Animal Welfare Committee of Berlin

(LAGeSo T-0124/08) and the permission of Biodonostia Institute Animal Care Committee. Wistar rats (FEM, Charité) were housed in specific pathogen-free barrier areas of the Hospital Charité Institute in Berlin while the C57B/6 (Jackson Laboratory) mice were housed in the Biodonostia Institute. Mice were maintained under a 12-hour light/12 hour dark cycle at 22° C with controlled humidity and with food and water provided ad libitum and handled in compliance with the animal research regulations specified in the European Communities Directive [2010/63/EU].

Astrocyte primary cultures

Primary astrocytes cell culture was prepared from neonatal Wistar rats during the first postnatal day of life. After dissection and careful removal of the meninges, both hemispheres were dissociated mechanically [13]. Cells were resuspended in Dulbecco's Modified Eagle Medium (DMEM, Invitrogen, USA) supplemented with 20% fetal calf serum, 1% penicillin/streptavidin and subsequently seeded in T75 flasks coated with poly-L-lysine (200 μ g/ml, Sigma-Aldrich) and grown in a humidified incubator maintained at 37° C under >90% humidity and 5% CO₂. Medium was changed every 2-3 days. After 7-10 days, the cultures were shaken overnight to minimize oligodendroglia and microglia contamination. For purification, the remaining astrocyte monolayers were trypsinized and replated. Cells were seeded in 12-well plates containing approximately 15,000 cells for immunocytochemistry and ATP analysis; in 6-well plates containing approximately 500,000 cells for quantitative RT-PCR and 10⁶ cells/well for western blot analysis.

Oxygen-glucose deprivation (OGD)

The culture medium was replaced with deoxygenated and glucose-free DMEM. Then cells were transferred to a humidified anaerobic chamber filled with a gas mixture of 5% CO₂ and 95% N₂ at 37° C. Throughout the OGD period of 4 hours, an atmosphere of <0.5% oxygen was maintained. After the OGD period, the media was quickly replaced with glucose-containing (1g/L) DMEM. Some samples were collected and others were reintroduced into normoxic conditions for 20 hours of reperfusion/recovery. Control cells in DMEM containing 4.5g/L glucose were always kept in a normoxic incubator.

OGD was performed 24h after cells were seeded (1DIV) or 4 weeks after (30DIV) to simulate an *in vitro* model of aging astrocytes [20]. The medium of the aged cells was changed every 2-3 days. For assays with P38MAPK inhibitor, astrocytes were treated with 2 μ M PH-797804 (Selleckchem) added to the medium when

seeded and included in every medium change. Control group were astrocytes with and without treatment in the different stages but without OGD.

ATP detection

ATP concentration was detected with the use of the Luminescent ATP Detection Assay Kit (Abcam, Cambridge, UK) according to the manufactures guide lines. Briefly, cultured astrocytes were incubated with a detergent on a shaker for 5 minutes. After total cell lyses, substrate for luminescence detection was added and incubated for 5 minutes. Luminescence was analyzed by IVIS Lumina *in vivo* Imaging System (PerkinElmer).

Tissue immunofluorescence

After mice perfusion, coronal serial sections of 50 μm were collected via SM2010 R Sliding Microtome (Leica) from young (2 months-old) and aged (2 years old) C57BL/6 mice and selected brain sections were blocked with 10% donkey serum and 0.1% Triton X-100 in phosphate-buffered saline and incubated with anti-P-p38MAPK (1:200; rabbit, Cell Signalling) and anti-GFAP conjugated with Alexa Fluor 488 (1:500; mouse, Sigma) overnight at 4° C. Nuclei were stained with DAPI staining (Sigma). Images were acquired using a Leica confocal microscope.

Cell immunofluorescence

After removal of the medium, cells seeded on coverslips were fixed with 4% paraformaldehyde for 15 min for immunocytochemical procedure. The samples were pre-incubated with a blocking solution (10% goat serum and 0.1% Triton X-100 in phosphate-buffered saline) for 30 min and then incubated with a mouse monoclonal anti-GFAP (1:400, Sigma, USA) for 2 hrs at room temperature, followed by incubation with rabbit anti cleaved caspase-3 (Asp175) (1:400, cell signaling) at 4° C overnight. After washes, cells were incubated with the respective secondary antibodies. Finally, after three washings, the sections were mounted with Vectashield HardSet Mounting Medium with DAPI (Sigma). Images were acquired using a Leica confocal microscope.

Quantitative RT-PCR

RNA extraction was performed with acidic phenol/chloroform (peq-GOLD RNApure, PEQLAB Biotechnologie, Erlangen, Germany) following the manufacturer protocol, and 2 μg of RNA were reverse transcribed. Primers were used for insulin-like growth factors (*Igf*), neuronal growth factor (*Ngf*), glial

fibrillary acidic protein (*Gfap*), superoxide dismutase 2 (*Sod2*), glutamine synthetase (*Gs*), glutamate aspartate transporter (*Slc1a3*), catalytic subunit of glutamate cysteine ligase (*Gclc*), to analyse the cDNA and *Hprt* was used as housekeeping gene (see Table 1). The expressions of target genes were analyzed with the StepOnePlus™ qPCR System (Applied Biosystems, Life Technologies, Carlsbad, CA) according to the $2^{-\Delta\Delta\text{CT}}$ method [27].

Western blot analysis

Immunoblots were performed following standard procedures. Equal amounts of protein (20 μg) were separated on 15% SDS polyacrylamide gels and blotted onto nitrocellulose membranes (BioRad). Primary antibodies were P-p38MAPK (1:200; Cell Signalling), total GFAP (1:100, Santa Cruz), TNF α (1:200, PromoCell), cleaved caspase 3 (1:1000, Cell Signaling) and α -ACTININ (1:2000, Sigma), followed by appropriate secondary antibodies conjugated with horseradish peroxidase (DAKO). Detection was performed by chemiluminescence using ECL (Amersham).

Statistical analysis

All data are presented as mean \pm standard error of the mean (SEM). After the assessment of normality with Kolmogorov Smirnov test (KS), t student test was performed to compare the differences between the control and the OGD group. Group differences were studied by one-way ANOVA with Bonferroni-Dunn correction. Two-sided $P < 0.05$ was considered statistically significant. The statistical analysis of data was performed using GraphPad prism 5 software version 5.01 (GraphPad Software, Inc. CA, USA).

RESULTS

Aged astrocytes show increased GFAP reactivity, inflammation, and cell death as well as lower nutritive and anti-oxidative capacity

We determined the GFAP reactivity in the cortex and dentate gyrus (DG) of the hippocampus in young (2 month-old) and aged (≥ 2 year-old) C57BL/6 mice. Immunofluorescence showed an increased expression of GFAP with aging (Figure 1A). To verify the age dependent increase of GFAP in astrocytes, we cultured rat neonatal astrocytes and simulated an *in vitro* model of aging (30DIV). Increased GFAP expression *in vivo* was confirmed in primary astrocytes *in vitro*. Aged astrocytes showed increased GFAP expression (0.6 ± 0.13 vs 1.3 ± 0.15 ratio vs DAPI) compared to younger cells (1DIV) (Figure 1B).

Table 1. Primer used in qPCR.

Gene	Forward primer	Reverse primer	GenBank ID
<i>Cntf</i>	AACCTTGACTCAGTGGATGGTGTGA	AAGCCTGGAGTTCTCTTGGA	NM_013166.1
<i>Gclc</i>	GGAGGACAACATGAGGAAACG	GCTCTGGCAGTGTGAATCCA	NM_012815.2
<i>Gfap</i>	TCTGGACCAGCTTACTACCAACAG	TGGTTTCATCTTGAGCTTCTG	NM_017009.2
<i>Igf-1</i>	CGGACCAGAGACCCTTTGC	GCCTGTGGGCTTGTGAAGT	NM_001082479.1
<i>Ngf</i>	ACCCAAGCTCACCTCAGTGTCT	GACATTACGCTATGCACCTCAGAGT	NM_001277055.1
<i>Slc1a3</i>	CCCTGCCCATCACTTTCAAG	GCGGTCCCATCCATGTTAAT	NM_001289942.1
<i>Sod2</i>	GACCTACGTGAACAATCTGAACGT	AGGCTGAAGAGCAACCTGAGTT	NM_017051.2
<i>Hprt</i>	GGAAAGAACGTCTTGATTGTTGAA	CCAACACTTCGAGAGGTCCTTTT	NM_012583.2

Mitochondria generate the majority most of cellular ATP supply, which is essential for energy consuming cell processes [28]. To investigate the impact of aging on cellular energy supply we measured the ATP concentration in 3ODIV (old) astrocytes and 1DIV (young). As a result, ATP luminescence was reduced in aged astrocytes to almost half of the level found in young cells ($P < 0.01$) (Figure 1C).

Regulation of glutamate homeostasis is one of the major functions of astrocytes in the brain. The enzyme glutamine synthetase (GS) is needed to transform glutamate intracellularly into glutamine, which is stored in vesicles [29]. In order to determine the impact of aging on glutamate homeostasis, we analyzed gene expression of *Gs* and of *Slc1a3* (EAAT1 or GLAST) representing a glutamate transporter molecule highly abundant in these cells [30]. qPCR analysis revealed that expression of *Gs* and *Slc1a3* is drastically reduced in aged astrocytes (Figure 1D, 1E). These results demonstrate that aging alters the expression of genes relevant for glutamate homeostasis in astrocytes.

The astrocytes anti-oxidant defense system is regulated via the Nrf2-pathway [31], which orchestrates the expression of its target genes including glutathione synthetase [32]. To define the impact of aging on anti-oxidant gene expression in astrocytes, we determined levels of *Sod2* and of catalytic subunit of glutamate cysteine ligase (*Gclc*) gene expression. The expression levels of *Sod2* and *Gclc* were significantly decreased in aged cultures indicating impairment of the anti-oxidant system in old astrocytes (Figure 1F, 1G).

To further characterize the mechanism of cell alterations caused by aging, we determined the expression of various growth factors, inflammatory response related genes, and performed caspase-3a (CASP3A) immunofluorescence staining for apoptosis. Aged astrocytes displayed a reduction of *igf* and an increase of *il6* (Figure 1H, 1I) gene expression. The numbers of CASP3A positive astrocytes in aged cultures were increased as compared to young cultures (2.9 ± 0.5 %

vs. 3.9 ± 0.3 %, number of CASP3A⁺ cells expressed as a percentage of all DAPI cells ($p < 0.05$) (Figure 1J). Western blot analysis corroborated the results of elevated GFAP and CASP3A expression in old cells (Figure 1K).

Loss of function in response to OGD is more pronounced in aged than in young astrocytes: role of cellular ATP levels, nutritive properties, and anti-oxidative capacity

To investigate the impact of OGD in young and aged astrocytes *in vitro*, we performed 4h of OGD (Figure 2A), and determined GFAP immunoreactivity in young and aged astrocytes. Immunofluorescence showed a significant OGD-induced increase of GFAP expression in both groups after OGD, which was even more pronounced in aged astrocytes (Figure 2B). The staining of GFAP was elevated exclusively in aged astrocytes after reperfusion/recovery.

We also determined the effect of OGD on cellular energy supply in young and aged astrocytes by ATP assay (Figure 2C). Reduction of ATP dependent energy after OGD in young astrocytes was restored back to control levels after 20 hours of reperfusion/recovery. Notably, in aged astrocytes, the OGD-induced decline of energy supply was still largely reduced after the phase of recovery (Figure 2C). This highlights that OGD hits aged astrocytic energy supply in a persistent way, and that the capacity of recovery and cellular repair is strongly impaired in old as compared to young astrocytes.

We also characterized the effect of OGD on glutamate uptake, anti-oxidative capacity, growth factor regulation, inflammatory response and cell death in young and aged astrocytes. Aging caused a significant upregulation of *Gs*, *Slc1a3*, *Sod2*, *Gclc* immediately after OGD indicating astroglial activation and induction of the anti-oxidant defense system compared to young astrocytes (Figure 2D–2G). OGD increased *Igf-1* expression only in aged astrocytes but its expression was




Figure 1. Aged astrocytes have increased reactivity, inflammation and cell death as well as loss of nutritive and anti-oxidative capacity. (A) Representative immunofluorescence for GFAP (red) in cortex and DG of young (2 month-old) and aged (over 24 month-old) C57BL/6J mice (n=2). (B) Representative immunofluorescence and the quantification for GFAP positive cells in 1DIV (young) and 30DIV (old) primary astrocytes cell culture derived from neonatal Wistar (n=6). (C) ATP luminescence levels of young and old primary astrocytes cultures (ratio compared to the young group) (n=6). (D–I) Expression of *gs*, *slc1a3*, *sod2*, *gclc*, *igf-1* and *il-6* in young and old primary astrocytes cultures (n=6). (J) Representative immunofluorescence of CASPASE 3A and co-staining of CASPASE 3A (red) with GFAP (green) together with DAPI (blue). Quantification for CASPASE 3A positive cells and in 1DIV (young) and 30DIV (old) primary astrocytes cell culture derived from P1 Wistar rat pups (n=6). (K) Protein expression of CASPASE 3A and GFAP in 1DIV (young) and 30DIV (old) primary astrocytes cell culture. Results are expressed as the mean \pm SEM. Asterisks denote the significance levels when compared to the control group (** $p < 0.001$, ** $p < 0.01$ and * $p < 0.05$ versus controls, t-test).




Figure 2. OGD has a higher impact in nutritive loss in aged compared to young astrocytes, whereas aged astrocytes have better anti-oxidant systems. (A) Representative optical microphotograph of young and old primary astrocytes derived from neonatal Wistar rats after 4h of OGD and after 20h of recovery compared to the control group. (B) Representative immunofluorescence and quantification of GFAP positive cells (green) in 1DIV (young) and 30DIV (old) primary astrocyte cell cultures after 4h of OGD and after 20h of recovery compared to the age matched control groups without OGD (n=6). (C) ATP luminiscence levels in young and old primary astrocyte cultures after 4h of OGD and after 20h of recovery compared to control groups (n=4). (D–H) Expression of *gs*, *slc1a3*, *sod2*, *gclc* and *igf-1* in young and old astrocytes after 4h of OGD and after 20h of recovery compared to controls (n=6). (I) Protein expression of TNF α in young and old cultured astrocytes after 4h of OGD compared to controls. (J) Representative immunofluorescence and the quantification for CASPASE 3A positive cells in 1DIV (young) and 30DIV (old) astrocytes cell culture derived from neonatal Wistar after 4h of OGD and after 20h of recovery compared to the control group (n=6). Results are expressed as the mean \pm SEM. Asterisks denote the significance levels when compared to the control group (**p<0.01, ***p<0.001, *p<0.05 versus controls, t-test).

downregulated after reperfusion/recovery in both groups (Figure 2H). Finally, there was a significant upregulation of TNF α and CASPASE3A after OGD in young and aged astrocytes (Figure 2I, 2J).

p38MAPK activity increases in aged astrocytes and its expression is reduced with PH-797804

p38MAPK activity (P-p38MAPK) and its isoforms are increased in different cell types in aged mouse brain [33]. Indeed, p38MAPK is one of the most important kinases in inflammatory signaling and its activation has been identified in many neurodegenerative diseases [21]. We first studied the expression of p38MAPK in the cortex and DG of hippocampus in young (2 month-old) and aged (≥ 2 year-old) C57BL/6J mice. Immunofluorescence showed that P-p38MAPK expression was increased in over 2 year-old animals

(Figure 3A), supporting previous studies in the DG [33]. Next, we measured P-p38MAPK in cultured young and aged astrocytes and found that P-p38MAPK expression was strongly increased in aged compared to young astrocytes (Figure 3B). Finally, the addition of PH-797804 to aged astrocyte cultures lowered levels of P-P38MAPK and *Mapk14* (p38alpha) gene expression by 50% after OGD and after reperfusion in comparison to control cells (Figure 3C, 3D).

Pharmacological inhibition of p38 α in aged astrocytes prevents astroglial reactivity, inflammatory response and anti-oxidant defense system activation after OGD

Finally, we aimed to clarify whether p38 α inhibition could have beneficial effects in aged astrocytes after 4h of OGD (Figure 4A). We first detected that *Mapk14*




Figure 3. p38MAPK activity increases in aged astrocytes and its expression is reduced with PH-797804. (A) Representative immunofluorescence and quantification for phosphorylated p38MAPK (P-p38MAPK) (green) together with DAPI (blue) and GFAP (red) in the cortex and in the DG of young (2 month-old) and aged (over 24 month-old) C57BL/6J mice (n=2). (B) Immunoblot of P-p38MAPK in 1DIV (young) and 30DIV (old) primary astrocyte cultures derived from Wistar rat brains. (C) Immunoblot of P-p38MAPK in old astrocyte cultures with and without PH-797804 treatment. (D) MAPK14 gene expression in old astrocyte cultures treated with PH-797804 at different time points in comparison to the control groups (without treatment) (n=6). Results are expressed as the mean \pm SEM. Asterisks denote the significance levels when compared to the control group (***)p<0.001, (**p<0.01 and (*p<0.05 versus controls, t-test).




Figure 4. Pharmacological inhibition of p38 α in aged astrocytes prevents from astrocytes reactivity, inflammatory response and anti-oxidant defense system activation after OGD. (A) Representative optical microphotograph of old primary astrocytes derived from neonatal Wistar after 4h of OGD and treated with PH-797804. (B) MAPK14 gene expression in old astrocyte cultures after 4h of OGD (white), after 4h of OGD with PH-797804 treatment (blue column) in comparison to controls treated with PH-797804 (black) (n=6). (C) Immunoblot of GFAP, TNF α and P-p38MAPK in old primary astrocytes cell culture derived from neonatal Wistar in normoxia and after 4h of OGD and with or without the treatment, PH-797804. (D) Representative quantification of GFAP positive cells in old astrocyte cultures derived after 4h of OGD, after 4h of OGD with PH-797804 treatment, and in control cultures treated with PH-797804 (n=6). (E–K) Expression of *gs*, *slc1a3*, *sod2*, *gclc*, *ngf*, *igf-1* and *il-6* in old astrocyte after 4h of OGD, after 4h of OGD with PH-797804 treatment, in controls treated with PH-797804 (n=6). Results are expressed as the mean \pm SEM. Asterisks denote the significance levels when compared to the control group (**p<0.001, **p<0.01 and *p<0.05 versus controls, t-test).

(p38MAPK alpha) expression was increased after OGD in aged astrocyte compared to the control group (Figure 4B). Analysis of TNF α , GFAP and P-p38MAPK protein expression after OGD revealed that there was a reduction of all these proteins in aged astrocytes treated with PH-797804 after OGD (Figure 4C).

We corroborated astroglial inhibition with GFAP immunofluorescence in which astrocytes treated with the p38 α inhibitor expressed lower levels of astrogliosis compared to the untreated OGD group directly after the injury and also after recovery (Figure 4D).

To further characterize the effect of pharmacological inhibition of p38 α in aged astrocytes after OGD and after recovery, we characterize the effects on glutamate uptake, anti-oxidative capacity and growth factor response. Gene expression of selected genes of those processes, *Gs*, *Slc1a3*, *Gclc*, *Sod*, *Ngf* and *Igf-1* was reduced in p38 α inhibited astrocytes treated with PH-797804 immediately after OGD and also after recovery (Figure 4E–4J). Hence, pharmacological inhibition of p38 α attenuates the oxidative challenge in aged astrocytes caused by OGD.

Finally, we analyzed the inflammatory response of astrocytes with PH-797804 treatment after OGD: The increase in aged astrocytes after 20h of recovery was fully abolished after pharmacological p38 α inhibition (Figure 4K).

DISCUSSION

In the central nervous system (CNS), astrocytes are the central cell type to support neuronal survival under pathological circumstances [34]. Indeed, as astrocytes, but not neurons, shift in gene expression patterns with aging, it has been proposed that they may be a better indicator of age in the brain than neurons [35]. Our *in vitro* results showed that aging increases the reactivity and inflammatory response of primary rat astrocytes. Previous reports show that astrocytes show age-dependent inflammatory responses [36] and increased GFAP expression [37, 38], which has been characterized as a hallmark of brain aging [5]. Moreover, our results indicate that the anti-oxidant and cellular energy system of *in vitro* aged astrocytes is decreased compared to the young ones, hence facilitating increased cell death. A proper cellular anti-oxidative capacity is necessary for oxidative stress protection, and a decline in ATP levels may contribute to neurodegeneration as it is related to regulating neuronal activity through synaptic inhibition [39]. The challenge through sublethal energy failure in our *in vitro* experiments may therefore lead or contribute to exhaustion of the anti-oxidant system. Furthermore, we found decreased *Gs* and *Slc1a3* expression in aged

astrocytes, hence pointing towards perturbed glutamate homeostasis as a potential trigger of cellular toxicity [29]. As a consequence, it can be assumed that protection of neurons via the anti-oxidant system of astrocytes is impaired by aging [40]. As a nutritive factor, IGF1 is considered to be essential for neuronal survival and development [41]. Moreover, previous studies have found that a lack of IGF1 reduces brain size with a cellular loss in the neuronal population and myelination deficits [42], but little attention has been paid to astrocytes as a source of IGF1, specifically. Consequently, the *Igf1* reduction found in our aged astrocytes may represent an injurious event to brain cells that were previously hit by energy failure. Altogether, the complex changes including inflammatory response, astrogliosis, energy failure, decreased anti-oxidative capacity and elevated caspase-3 activity may contribute to an increased “*in vitro*” aged astrocyte cell death.

As mentioned, aging alone increases astroglial reactivity [43], which is an observation also supported by our results, but this glial response is exaggerated following ischemia-stroke in the brain, hence accelerating glial scar formation [4]. We found that responses of energy failure and inflammation were exaggerated following 4 hours of OGD in our *in vitro* aged astrocytes compared to the young astrocytes, and that these alterations were maintained during the phase of recovery. However, although aged astrocytes present decreased anti-oxidative capacity, after OGD we observed an upregulation of anti-oxidant system defense, growth factor release and glutamate uptake. Cellular defense in the brain involve endogenous protective enzymes such as SOD and glutathione (GSH), which are produced by astrocytes to protect neurons against oxidative stress. In previous studies it has been demonstrated that the cellular anti-oxidant system and GSH release is upregulated in astrocytes as a response to pathological circumstances, possibly as a defense mechanism to prevent cell apoptosis [44]. Hence, the increase of cellular anti-oxidant defense system, *Gs* and *Slc1a3* expression and growth factor expression may represent a cellular attempt of compensation in aged astrocytes in response to astrogliosis, inflammation and CASPASE 3 activation induced by OGD.

Our results show enhanced p38MAPK activity with aging in astrocytes, both in astrocyte cultures *in vitro* and in the cortex and dentate gyrus (hippocampus) *in vivo*, which complement previous results obtained in neurogenic niches (SVZ) *in vivo* and in neurospheres derived from aged mice [45]. Elevated p38MAPK activity has also been identified in neurodegenerative disease and in response to brain injury such as brain stroke [46], and its pharmacological inhibition ameliorated symptoms of neurodegenerative diseases

and was protective against ischemia [21, 25]. We show that *Mapk14* expression was increased in aged astrocytes directly after OGD and that pharmacological p38 α inhibition in aged astrocytes after OGD restored *Mapk14* expression back to control levels. Moreover, the P38MAPK pathway has been described as inflammatory mediator in the CNS [33]. In our results, the inactivation of p38 α in aged astrocyte cultures treated by PH-797804 attenuated astroglial activation and inflammation that occur after OGD. We also found that PH-797804 treatment in aged astrocytes can prevent OGD-induced changes of growth factors *igf* and *ngf*, of the free radical clearance factors *sod2* and *gclc*, and of the glutamate metabolism system *gs* that were increased after OGD. The impaired production of these factors has previously been reported to alter the supportive properties of astrocyte [13, 47].

In summary, our experiments with OGD highlight an impairment of various astroglial functions that are important for the support of neuronal development and for protection of the brain both in young and aged astrocytes cell culture. Inhibition of p38 α in aged *in vitro* astrocytes may provide effective protection reducing astrogliosis and inflammation that occurs in aged astrocyte after OGD event.

Ethics approval

All procedures were approved by the state animal welfare authorities Berlin, Germany (LAGeSo T-0124/08) and followed institutional guidelines.

AUTHOR CONTRIBUTIONS

MR, AE and TS performed all the experiments the acquisition of data, analysis and interpretation of data. SE and LM-C contributed to the interpretation of data and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. CB and AM contributed to the final approval of the version to be published and acquisition of funding. TS directed the project, contributed to conception, design and analysis and interpretation of data and has been involved in drafting the manuscript and revising it critically for important intellectual content.

ACKNOWLEDGMENTS

We thank Mrs. Ruth Herrmann for help with immunohistochemistry and Elisa techniques.

CONFLICTS OF INTEREST

None of the authors have any conflicts of interest.

FUNDING

Deutsche Forschungsgemeinschaft (SCHE 2078/2-1). Förderverein für frühgeborene Kinder an der Charité e.V. Basque Government Postdoc (2017_1_0095).

REFERENCES

1. Sarikaya H, Steinlin M. Cerebellar stroke in adults and children. *Handb Clin Neurol*. 2018; 155:301–12. <https://doi.org/10.1016/B978-0-444-64189-2.00020-2> PMID:[29891068](https://pubmed.ncbi.nlm.nih.gov/29891068/)
2. Wallace AN, Kansagra AP, McEachern J, Moran CJ, Cross DT 3rd, Derdeyn CP. Evolution of endovascular stroke therapies and devices. *Expert Rev Med Devices*. 2016; 13:263–70. <https://doi.org/10.1586/17434440.2016.1143772> PMID:[26781520](https://pubmed.ncbi.nlm.nih.gov/26781520/)
3. Chen RL, Balami JS, Esiri MM, Chen LK, Buchan AM. Ischemic stroke in the elderly: an overview of evidence. *Nat Rev Neurol*. 2010; 6:256–65. <https://doi.org/10.1038/nrneurol.2010.36> PMID:[20368741](https://pubmed.ncbi.nlm.nih.gov/20368741/)
4. Badan I, Buchhold B, Hamm A, Gratz M, Walker LC, Platt D, Kessler C, Popa-Wagner A. Accelerated glial reactivity to stroke in aged rats correlates with reduced functional recovery. *J Cereb Blood Flow Metab*. 2003; 23:845–54. <https://doi.org/10.1097/01.WCB.0000071883.63724.A7> PMID:[12843788](https://pubmed.ncbi.nlm.nih.gov/12843788/)
5. Jiang T, Cadenas E. Astrocytic metabolic and inflammatory changes as a function of age. *Aging Cell*. 2014; 13:1059–67. <https://doi.org/10.1111/acer.12268> PMID:[25233945](https://pubmed.ncbi.nlm.nih.gov/25233945/)
6. Back SA, Craig A, Kayton RJ, Luo NL, Meshul CK, Allcock N, Fern R. Hypoxia-ischemia preferentially triggers glutamate depletion from oligodendroglia and axons in perinatal cerebral white matter. *J Cereb Blood Flow Metab*. 2007; 27:334–47. <https://doi.org/10.1038/sj.jcbfm.9600344> PMID:[16757980](https://pubmed.ncbi.nlm.nih.gov/16757980/)
7. Lipton SA. Molecular mechanisms of trauma-induced neuronal degeneration. *Curr Opin Neurol Neurosurg*. 1993; 6:588–96. PMID:[8400474](https://pubmed.ncbi.nlm.nih.gov/8400474/)
8. Fernández-López D, Faustino J, Daneman R, Zhou L, Lee SY, Derugin N, Wendland MF, Vexler ZS. Blood-brain barrier permeability is increased after acute adult stroke but not neonatal stroke in the rat. *J Neurosci*. 2012; 32:9588–600. <https://doi.org/10.1523/JNEUROSCI.5977-11.2012> PMID:[22787045](https://pubmed.ncbi.nlm.nih.gov/22787045/)

9. Sommer CJ. Ischemic stroke: experimental models and reality. *Acta Neuropathol.* 2017; 133:245–61.
<https://doi.org/10.1007/s00401-017-1667-0>
PMID:[28064357](https://pubmed.ncbi.nlm.nih.gov/28064357/)
10. Schmitz T, Endesfelder S, Chew LJ, Zaak I, Bühner C. Minocycline protects oligodendroglial precursor cells against injury caused by oxygen-glucose deprivation. *J Neurosci Res.* 2012; 90:933–44.
<https://doi.org/10.1002/jnr.22824> PMID:[22253205](https://pubmed.ncbi.nlm.nih.gov/22253205/)
11. Goux E, Buisson A, Nieoullon A, Kerkerian-Le Goff L, Tauskela JS, Blondeau N, Had-Aissouni L. Oxygen glucose deprivation-induced astrocyte dysfunction provokes neuronal death through oxidative stress. *Pharmacol Res.* 2014; 87:8–17.
<https://doi.org/10.1016/j.phrs.2014.06.002>
PMID:[24928737](https://pubmed.ncbi.nlm.nih.gov/24928737/)
12. Clemente D, Ortega MC, Melero-Jerez C, de Castro F. The effect of glia-glia interactions on oligodendrocyte precursor cell biology during development and in demyelinating diseases. *Front Cell Neurosci.* 2013; 7:268.
<https://doi.org/10.3389/fncel.2013.00268>
PMID:[24391545](https://pubmed.ncbi.nlm.nih.gov/24391545/)
13. Scheuer T, Brockmüller V, Blanco Knowlton M, Weitkamp JH, Ruhwedel T, Mueller S, Endesfelder S, Bühner C, Schmitz T. Oligodendroglial maldevelopment in the cerebellum after postnatal hyperoxia and its prevention by minocycline. *Glia.* 2015; 63:1825–39.
<https://doi.org/10.1002/glia.22847> PMID:[25964099](https://pubmed.ncbi.nlm.nih.gov/25964099/)
14. Bucher F, Stahl A, Agostini HT, Martin G. Hyperoxia causes reduced density of retinal astrocytes in the central avascular zone in the mouse model of oxygen-induced retinopathy. *Mol Cell Neurosci.* 2013; 56:225–33.
<https://doi.org/10.1016/j.mcn.2013.06.001>
PMID:[23756201](https://pubmed.ncbi.nlm.nih.gov/23756201/)
15. Miyamoto N, Maki T, Shindo A, Liang AC, Maeda M, Egawa N, Itoh K, Lo EK, Lok J, Ihara M, Arai K. Astrocytes promote oligodendrogenesis after white matter damage via brain-derived neurotrophic factor. *J Neurosci.* 2015; 35:14002–08.
<https://doi.org/10.1523/JNEUROSCI.1592-15.2015>
PMID:[26468200](https://pubmed.ncbi.nlm.nih.gov/26468200/)
16. Xu L, Emery JF, Ouyang YB, Voloboueva LA, Giffard RG. Astrocyte targeted overexpression of Hsp72 or SOD2 reduces neuronal vulnerability to forebrain ischemia. *Glia.* 2010; 58:1042–49.
<https://doi.org/10.1002/glia.20985>
PMID:[20235222](https://pubmed.ncbi.nlm.nih.gov/20235222/)
17. Zhang R, Huang Q, Zou L, Cao X, Huang H, Chu X. Beneficial effects of deferoxamine against astrocyte death induced by modified oxygen glucose deprivation. *Brain Res.* 2014; 1583:23–33.
<https://doi.org/10.1016/j.brainres.2014.08.016>
PMID:[25152469](https://pubmed.ncbi.nlm.nih.gov/25152469/)
18. Revuelta M, Arteaga O, Alvarez A, Martinez-Ibargüen A, Hilario E. Characterization of gene expression in the rat brainstem after neonatal hypoxic-ischemic injury and antioxidant treatment. *Mol Neurobiol.* 2017; 54:1129–43.
<https://doi.org/10.1007/s12035-016-9724-6>
PMID:[26809461](https://pubmed.ncbi.nlm.nih.gov/26809461/)
19. Zhao H, Liu Y, Cheng L, Liu B, Zhang W, Guo YJ, Nie L. Mesencephalic astrocyte-derived neurotrophic factor inhibits oxygen-glucose deprivation-induced cell damage and inflammation by suppressing endoplasmic reticulum stress in rat primary astrocytes. *J Mol Neurosci.* 2013; 51:671–78.
<https://doi.org/10.1007/s12031-013-0042-4>
PMID:[23760988](https://pubmed.ncbi.nlm.nih.gov/23760988/)
20. Caldeira C, Oliveira AF, Cunha C, Vaz AR, Falcão AS, Fernandes A, Brites D. Microglia change from a reactive to an age-like phenotype with the time in culture. *Front Cell Neurosci.* 2014; 8:152.
<https://doi.org/10.3389/fncel.2014.00152>
PMID:[24917789](https://pubmed.ncbi.nlm.nih.gov/24917789/)
21. Colié S, Sarroca S, Palenzuela R, Garcia I, Matheu A, Corpas R, Dotti CG, Esteban JA, Sanfeliu C, Nebreda AR. Neuronal p38 α mediates synaptic and cognitive dysfunction in an Alzheimer's mouse model by controlling β -amyloid production. *Sci Rep.* 2017; 7:45306.
<https://doi.org/10.1038/srep45306>
PMID:[28361984](https://pubmed.ncbi.nlm.nih.gov/28361984/)
22. Krupinski J, Slevin M, Marti E, Catena E, Rubio F, Gaffney J. Time-course phosphorylation of the mitogen activated protein (MAP) kinase group of signalling proteins and related molecules following middle cerebral artery occlusion (MCAO) in rats. *Neuropathol Appl Neurobiol.* 2003; 29:144–58.
<https://doi.org/10.1046/j.1365-2990.2003.00454.x>
PMID:[12662322](https://pubmed.ncbi.nlm.nih.gov/12662322/)
23. Cuadrado A, Nebreda AR. Mechanisms and functions of p38 MAPK signalling. *Biochem J.* 2010; 429:403–17.
<https://doi.org/10.1042/BJ20100323> PMID:[20626350](https://pubmed.ncbi.nlm.nih.gov/20626350/)
24. Roy Choudhury G, Ryou MG, Poteet E, Wen Y, He R, Sun F, Yuan F, Jin K, Yang SH. Involvement of p38 MAPK in reactive astrogliosis induced by ischemic stroke. *Brain Res.* 2014; 1551:45–58.
<https://doi.org/10.1016/j.brainres.2014.01.013>
PMID:[24440774](https://pubmed.ncbi.nlm.nih.gov/24440774/)
25. Roy SM, Grum-Tokars VL, Schavocky JP, Saeed F, Staniszewski A, Teich AF, Arancio O, Bachstetter AD, Webster SJ, Van Eldik LJ, Minasov G, Anderson WF, Pelletier JC, Watterson DM. Targeting human central

- nervous system protein kinases: an isoform selective p38 α MAPK inhibitor that attenuates disease progression in Alzheimer's disease mouse models. *ACS Chem Neurosci*. 2015; 6:666–80.
<https://doi.org/10.1021/acschemneuro.5b00002>
PMID:25676389
26. Barone FC, Irving EA, Ray AM, Lee JC, Kassis S, Kumar S, Badger AM, Legos JJ, Erhardt JA, Ohlstein EH, Hunter AJ, Harrison DC, Philpott K, et al. Inhibition of p38 mitogen-activated protein kinase provides neuroprotection in cerebral focal ischemia. *Med Res Rev*. 2001; 21:129–45.
[https://doi.org/10.1002/1098-1128\(200103\)21:2<129::aid-med1003>3.0.co;2-h](https://doi.org/10.1002/1098-1128(200103)21:2<129::aid-med1003>3.0.co;2-h)
PMID:11223862
 27. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2⁻(Delta Delta C(T)) method. *Methods*. 2001; 25:402–08.
<https://doi.org/10.1006/meth.2001.1262>
PMID:11846609
 28. Consolini AE, Ragone MI, Bonazzola P, Colareda GA. Mitochondrial bioenergetics during ischemia and reperfusion. *Adv Exp Med Biol*. 2017; 982:141–67.
https://doi.org/10.1007/978-3-319-55330-6_8
PMID:28551786
 29. Lin CH, You JR, Wei KC, Gean PW. Stimulating ERK/PI3K/NF κ B signaling pathways upon activation of mGluR2/3 restores OGD-induced impairment in glutamate clearance in astrocytes. *Eur J Neurosci*. 2014; 39:83–96.
<https://doi.org/10.1111/ejn.12383>
PMID:24206109
 30. Llorente IL, Landucci E, Pellegrini-Giampietro DE, Fernández-López A. Glutamate receptor and transporter modifications in rat organotypic hippocampal slice cultures exposed to oxygen-glucose deprivation: the contribution of cyclooxygenase-2. *Neuroscience*. 2015; 292:118–28.
<https://doi.org/10.1016/j.neuroscience.2015.02.040>
PMID:25732138
 31. Wu L, Li HH, Wu Q, Miao S, Liu ZJ, Wu P, Ye DY. Lipoxin A4 activates Nrf2 pathway and ameliorates cell damage in cultured cortical astrocytes exposed to oxygen-glucose deprivation/reperfusion insults. *J Mol Neurosci*. 2015; 56:848–57.
<https://doi.org/10.1007/s12031-015-0525-6>
PMID:25702137
 32. Li L, Yang N, Nin L, Zhao Z, Chen L, Yu J, Jiang Z, Zhong Z, Zeng D, Qi H, Xu X. Chinese herbal medicine formula tao hong si wu decoction protects against cerebral ischemia-reperfusion injury via PI3K/Akt and the Nrf2 signaling pathway. *J Nat Med*. 2015; 69:76–85.
<https://doi.org/10.1007/s11418-014-0865-5>
PMID:25149059
 33. Moreno-Cugnon L, Revuelta M, Arrizabalaga O, Colie S, Moreno-Valladares M, Jimenez-Blasco D, Gil-Bea F, Llarena I, Bolaños JP, Nebreda AR, Matheu A. Neuronal p38 α mediates age-associated neural stem cell exhaustion and cognitive decline. *Aging Cell*. 2019; 18:e13044.
<https://doi.org/10.1111/accel.13044> PMID:31560167
 34. Barnett SC, Linington C. Myelination: do astrocytes play a role? *Neuroscientist*. 2013; 19:442–50.
<https://doi.org/10.1177/1073858412465655>
PMID:23131748
 35. Soreq L, Rose J, Soreq E, Hardy J, Trabzuni D, Cookson MR, Smith C, Ryten M, Patani R, Ule J, and UK Brain Expression Consortium, and North American Brain Expression Consortium. Major shifts in glial regional identity are a transcriptional hallmark of human brain aging. *Cell Rep*. 2017; 18:557–70.
<https://doi.org/10.1016/j.celrep.2016.12.011>
PMID:28076797
 36. Campuzano O, Castillo-Ruiz MM, Acarin L, Castellano B, Gonzalez B. Increased levels of proinflammatory cytokines in the aged rat brain attenuate injury-induced cytokine response after excitotoxic damage. *J Neurosci Res*. 2009; 87:2484–97.
<https://doi.org/10.1002/jnr.22074> PMID:19326443
 37. Cohen J, Torres C. Astrocyte senescence: evidence and significance. *Aging Cell*. 2019; 18:e12937.
<https://doi.org/10.1111/accel.12937> PMID:30815970
 38. Clarke LE, Liddelow SA, Chakraborty C, Münch AE, Heiman M, Barres BA. Normal aging induces A1-like astrocyte reactivity. *Proc Natl Acad Sci USA*. 2018; 115:E1896–905.
<https://doi.org/10.1073/pnas.1800165115>
PMID:29437957
 39. Lalo U, Rasooli-Nejad S, Pankratov Y. Exocytosis of gliotransmitters from cortical astrocytes: implications for synaptic plasticity and aging. *Biochem Soc Trans*. 2014; 42:1275–81.
<https://doi.org/10.1042/BST20140163> PMID:25233403
 40. Ishii T, Takanashi Y, Sugita K, Miyazawa M, Yanagihara R, Yasuda K, Onouchi H, Kawabe N, Nakata M, Yamamoto Y, Hartman PS, Ishii N. Endogenous reactive oxygen species cause astrocyte defects and neuronal dysfunctions in the hippocampus: a new model for aging brain. *Aging Cell*. 2017; 16:39–51.
<https://doi.org/10.1111/accel.12523> PMID:27623715
 41. Wang J, Tang Y, Zhang W, Zhao H, Wang R, Yan Y, Xu L, Li P. Insulin-like growth factor-1 secreted by brain microvascular endothelial cells attenuates neuron injury upon ischemia. *FEBS J*. 2013; 280:3658–68.

<https://doi.org/10.1111/febs.12359>

PMID:[23721666](https://pubmed.ncbi.nlm.nih.gov/23721666/)

42. Seto D, Zheng WH, McNicoll A, Collier B, Quirion R, Kar S. Insulin-like growth factor-I inhibits endogenous acetylcholine release from the rat hippocampal formation: possible involvement of GABA in mediating the effects. *Neuroscience*. 2002; 115:603–12.
[https://doi.org/10.1016/s0306-4522\(02\)00450-5](https://doi.org/10.1016/s0306-4522(02)00450-5)
PMID:[12421625](https://pubmed.ncbi.nlm.nih.gov/12421625/)
43. Lively S, Moxon-Emre I, Schlichter LC. SC1/hevin and reactive gliosis after transient ischemic stroke in young and aged rats. *J Neuropathol Exp Neurol*. 2011; 70:913–29.
<https://doi.org/10.1097/NEN.0b013e318231151e>
PMID:[21937915](https://pubmed.ncbi.nlm.nih.gov/21937915/)
44. Alfieri A, Srivastava S, Siow RC, Modo M, Fraser PA, Mann GE. Targeting the Nrf2-Keap1 antioxidant defence pathway for neurovascular protection in stroke. *J Physiol*. 2011; 589:4125–36.
45. Moreno-Cugnion L, Arrizabalaga O, Llarena I, Matheu A. Elevated p38MAPK activity promotes neural stem cell aging. *Aging (Albany NY)*. 2020; 12:6030–36.
<https://doi.org/10.18632/aging.102994>
PMID:[32243258](https://pubmed.ncbi.nlm.nih.gov/32243258/)
46. Irving EA, Barone FC, Reith AD, Hadingham SJ, Parsons AA. Differential activation of MAPK/ERK and p38/SAPK in neurones and glia following focal cerebral ischaemia in the rat. *Brain Res Mol Brain Res*. 2000; 77:65–75.
[https://doi.org/10.1016/s0169-328x\(00\)00043-7](https://doi.org/10.1016/s0169-328x(00)00043-7)
PMID:[10814833](https://pubmed.ncbi.nlm.nih.gov/10814833/)
47. Revuelta M, EliceGUI A, Moreno-Cugnion L, Bührer C, Matheu A, Schmitz T. Ischemic stroke in neonatal and adult astrocytes. *Mech Ageing Dev*. 2019; 183:111147.
<https://doi.org/10.1016/j.mad.2019.111147>
PMID:[31493435](https://pubmed.ncbi.nlm.nih.gov/31493435/)