

# Hyperbaric oxygen therapy induces transcriptome changes in elderly: a prospective trial

Amir Hadanny<sup>1,2,3,4</sup>, Relly Forer<sup>5</sup>, Dina Volodarsky<sup>5</sup>, Malka Daniel-Kotovsky<sup>1</sup>, Merav Catalogna<sup>1</sup>, Yonatan Zemel<sup>1,4</sup>, Yair Bechor<sup>1,4</sup>, Shai Efrati<sup>1,2,4,6</sup>

<sup>1</sup>The Sagol Center for Hyperbaric Medicine and Research, Shamir (Assaf-Harofeh) Medical Center, Zerifin, Israel

<sup>2</sup>Sackler School of Medicine, Tel-Aviv University, Tel-Aviv, Israel

<sup>3</sup>Bar Ilan University, Ramat-Gan, Israel

<sup>4</sup>Aviv Scientific LTD, Bnei-Brak, Israel

<sup>5</sup>Dyn Diagnostics, Zerifin, Israel

<sup>6</sup>Sagol School of Neuroscience, Tel-Aviv University, Tel-Aviv, Israel

**Correspondence to:** Amir Hadanny; email: [amir.had@gmail.com](mailto:amir.had@gmail.com), <https://orcid.org/0000-0001-9028-7268>

**Keywords:** transcriptome, aging, hyperbaric oxygen, HBOT, RNA

**Received:** November 23, 2020

**Accepted:** November 11, 2021

**Published:** November 24, 2021

**Copyright:** © 2021 Hadanny et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/3.0/) (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## ABSTRACT

**Introduction:** Aging is characterized by the progressive loss of physiological capacity. Changes in gene expression can alter activity in defined age-related molecular pathways leading to cellular aging and increased aging disease susceptibility. The aim of the current study was to evaluate whether hyperbaric oxygen therapy (HBOT) affects gene expression in normal, non-pathological, aging adults.

**Methods:** Thirty-five healthy independently living adults, aged 64 and older, were enrolled to receive 60 daily HBOT exposures. Whole blood samples were collected at baseline, at the 30th and 60th HBOT session, and 1–2 weeks following the last session. Differential gene expression analysis was performed.

**Results:** Following 60 sessions of HBOT, 1342 genes and 570 genes were differently up- and downregulated (1912 total), respectively ( $p < 0.01$  FDR), compared to baseline. Out of which, five genes were downregulated with a  $>1.5$ -fold change: ABCA13 (FC =  $-2.28$ ), DNAJ6 (FC =  $-2.16$ ), HBG2 (FC =  $-1.56$ ), PDXDC1 (FC =  $-1.53$ ), RANBP17 (FC =  $-1.75$ ). Two weeks post-HBOT, ABCA13 expression was significantly downregulated with a  $>1.5$ fold change (FC =  $-1.54$ ,  $p = 0.008$ ).

**In conclusion,** for the first time in humans, the study provides direct evidence of HBOT is associated with transcriptome changes in whole-blood samples. Our results demonstrate significant changes in gene expression of normal aging population.

## INTRODUCTION

Aging can be defined by the gradual decline of physiological capacities, resulting in impaired functions and susceptibility for diseases and death. This biological deterioration is considered a major risk factor for Alzheimer's disease, cardiovascular disease, diabetes mellitus, cancer among others. The aging rate is partially determined, by genetics and other evolutionary conserved biochemical processes [1].

Gene expression (whole transcriptome) interrogation is widely used to explore differences between individual young and old, and diseased and healthy populations [2–4]. Significant transcriptional changes related to aging have been suggested with respect to inflammation, oxidative stress, mitochondrial and lysosomal degradation pathways associated genes [3]. Moreover, these gene expression changes can alter activity in defined age-related molecular pathways leading to cellular aging and increased susceptibility to aging diseases.

Hyperbaric oxygen therapy (HBOT) utilizes pure oxygen (100%) at high barometric pressure (>at least over one absolute atmospheres (ATA)) to enhance the body's tissues oxygen content. Specific HBOT protocols which utilize repeated intermittent hyperoxic exposures induce physiological effects which are classically associated with hypoxia, only using the hyperoxic environment instead. This has been previously referred as the hyperoxic-hypoxic paradox [5–8]. Clinically, HBOT has been used for non-healing ischemic wounds and post-radiation injuries by promoting angiogenesis and wound healing [9]. In our previous study, our group provided evidence HBOT HBOT can induce significant cognitive improvement in the healthy aging population, mediated by cerebral blood flow changes [10]. This indicates that HBOT's pleiotropic regenerative effects are mediated by activating various genetic pathways. Previous studies evaluated HBOT's effects on isolated cell cultures and identified different genes sensitive to pressure and oxygen changes including inflammatory, growth, repair, angiogenesis, tumor suppressors, stress, cellular stress and apoptosis associated genes [11–14]. So far, HBOT effects on gene expression *in-vivo* has yet to be explored in clinical studies.

The aim of the current study was to evaluate whether HBOT affects gene expression in the normal aging population (excluding pathological aging).

## METHODS

### Subjects

Thirty-five aging independent adults good functional and cognitive status over 64 years old were enrolled. The clinical study was performed in the Shamir (Assaf-Harofeh) Medical Center, Israel between 2016–2020. All included participants ruled out any history of cardiac or cerebrovascular ischemia for at least one year prior to inclusion. Patients with previous HBOT exposure in the last three months prior to inclusion, known malignancy in the year prior to inclusion, pathological cognitive decline, uncontrolled diabetes mellitus (HbA1C>8, fasting glucose>200), severe chronic renal failure (GFR <30), immunosuppressants, MRI contraindications or active smoking or pulmonary diseases were excluded.

### Study design

The study protocol was approved by the Shamir Medical Center's Institutional Review Board. All patients signed an informed consent followed by baseline evaluations, HBOT protocol and post therapy evaluations. Measurement points were evaluated at

baseline, 2nd baseline control following 60–100 days, last (60th) HBOT session and 7–14 days following the last HBOT session.

The study cohort included only HBOT treated patients, who are part of a larger cohort of normal ageing population studied at the Shamir Medical Center, Israel (NCT02790541 [10]).

### Interventions

The HBOT protocol was provided using a multiplace chamber (Starmed-2700, HAUX, Germany). The protocol included 60 daily sessions, during a three-month period with five sessions per week. Each session comprised of 100% oxygen provided by mask at 2ATA for 90 minutes and intermittent air breaks for five minutes every 20 minutes of oxygen. Compression and decompression rates were controlled at 1 meter/minute. During the study, lifestyle and diet changes and medication adjustments were not allowed.

### Blood samples

Whole blood samples were collected into Tempus™ blood RNA tubes containing a stabilizing reagent using a standard technique following overnight fasting, at baseline, at the second control following 60–100 days, the day of the last HBOT session (60th session) and 10–14 days following HBOT protocol.

### RNA extraction

RNA was extracted using a Tempus™ Spin RNA Isolation Kit (LifeTechnologies, Warrington, UK). Spectrophotometer (NanoDrop 1000, Thermo Fisher Scientific, Waltham, MA, United States) was used to quantify total RNA samples followed by integrity confirmation with the Agilent Bioanalyzer 2100 with an RNA Nano-Chip Kit (Agilent Technologies, Waldbronn, Germany). One hundred nanograms of RNA were used for the Clariom S assay (Affymetrix) following manufacturer's standardized protocol. The Clariom assay interrogates over 22,000 annotated genes, each consisting of 6–10 probes exactly matching the target transcript sequence.

In brief, 100 ng total RNA and dNTP-T7 random primers were used for reverse transcription (RT). Amplified cRNA was made using Second strand synthesis and *in vitro* transcription amplified cRNA. Subsequent reverse transcription produced cDNA, followed by fragmentation. Fragments were labeled with biotin for hybridization to the replication and transcription activator (RTA). Array processing was done using an Affymetrix GeneChip® Hybridization,

Wash, and Stain kit. Initially, 5.2 µg biotin-labeled fragmented ss-cDNA was added to 1× hybridization buffer containing oligo B2, hybridization controls, and DMSO. Following incubation at 95°C and 45°C for 5 minutes each, 200 µL hybridization mix was loaded onto each array. Arrays were incubated for 16th in a GeneChip 645 hybridization oven (Affymetrix) at 45°C while rotating at 60 rpm. Wash/stain procedures were performed on an Affymetrix 450 Fluidics Station using manufacturer specified instrument protocol settings. To remove the unbound sample, arrays were washed with non-stringent wash-Buffer A. GeneChips® were then stained 10 min in Stain Cocktail 1. Excess stain was removed by a subsequent wash in Buffer A. Arrays were incubated 10 min with Stain Cocktail 2 followed by 10 min incubation with Stain Cocktail 1 for signal amplification. GeneChips were washed with Buffer B then filled with Array Holding Buffer prior to removal from the fluidics station and scanned using the GeneArray® 3000 scanner (see Affymetrix target hybridization protocol).

### Transcriptome analysis

The Affymetrix Transcriptome Analysis Console software (TAC) ver. 4.0.1.36 (ThermoFisher Scientific) was used for primary data analysis. Data were normalized using the TAC Robust Multiarray Average module. The Limma Bioconductor package was used for differential expression analysis using an empirical Bayesian correction approach. Gene expression data were log-transformed.

Both baseline measurements were combined as one factor to exclude non-HBOT related changes. A repeated measures one-way ANOVA was used to compare differentially expressed genes (DEG) FC between the 60th session measurement and baseline, post-treatment measurement and baseline. Then a Benjamini-Hochberg method [15] was applied on the calculated *p*-values to correct for false discovery rate. A change was considered significant if met the criterion *P* < 0.01 following a false discovery rate correction.

Functional terms were extracted using the Gene Ontology (GO) terms and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways. The Expander software was used for both GO and KEGG pathway enrichment analyses [16]. False discovery rate (FDR) below 0.05 threshold was used for statistically significant KEGG pathways and GO terms.

### Quantitative real-time PCR (qPCR)

In order to confirm microarray results RT qPCR reactions were performed. The RT reaction process of mRNA was

done with oligo dT primers. The SYBR Green PCR kit (Toyobo, Osaka, Japan) was used to perform qPCR on LightCycler480 real-time PCR system (Roche Diagnostics). HPRT and ABCA13 were the main reference targets of mRNA. The following primers were used: HPRT forward 5' TTGTTGTAGGATATGCCCTTGA 3' and HPRT reverse 5' GCGATGTCAATAGGACTCCAG 3', ABCA13 forward 5' GCTTTCTGTATCCTAGTTCCTTCTGT 3' and ABCA13 rev 5' GATGTACTCTCGCCTCCTAGAT 3'.

### Statistical analysis

Unless otherwise specified (see above for transcriptome analysis), demographic continuous data are expressed as means ± standard-deviation. The Kolmogorov-Smirnov test was used to determine normal distribution for all variables.

Demographics categorical data are expressed in numbers and percentages. Univariate analyses were performed using either chi-square or Fisher's exact tests, as appropriate. Correlation was performed using Pearson's correlation coefficient.

## RESULTS

Thirty-five individuals were assigned to HBOT. Five patients were excluded since they did not complete baseline assessments. All other 30 patients completed baseline evaluations the HBOT protocol. Three blood samples did not include RNA tubes and were excluded from analysis (Figure 1). The average age was 70.39 ± 3.72 and 51.9% were males. The baseline characteristics and comparison of the cohorts following exclusion of the patients are in Table 1.

Following 60 sessions of HBOT, 1342 genes and 570 genes were differently up- and downregulated (1912 total), respectively (*p* < 0.01 FDR) compared to baseline (Figure 2). Out of which, five genes were downregulated with a >1.5-fold change: ABCA13 (FC = -2.28), DNAJ6 (FC = -2.16), HBG2 (FC = -1.56), PDXDC1 (FC = -1.53), RANBP17 (FC = -1.75) (Table 2). The full list of obtained DEGs are in Supplementary Table 1.

Two weeks post-HBOT, 11 genes and 8 genes were differently up- and downregulated (19 total), respectively (*p* < 0.01, FDR) compared to baseline (Figure 3). Out of which, ABCA13 expression was significantly downregulated with a >1.5-fold change (FC = -1.54, *p* = 0.008). The full list of obtained DEGs are in Supplementary Table 2.

The overlap of the two timepoints (Last HBOT session vs baseline, and two weeks post last HBOT sessions vs

baseline) consisted of nine differentially expressed genes (DEGs), including *ABCA13*, *RNF13*, *BOC*, *ARL2*, *MRPS52*, *ATP5J2*, *C20orf27*, *SYNGAP1*, *RHOBTB2*. However the only gene with more than 1.5-fold change was *ABCA13* (Table 3).

To validate this finding, RT-qPCR was performed for *ABCA13* gene on all available samples (16/27) with an average fold change of  $-3.89 \pm 4.45$ , where in 12/16 the absolute fold change was higher than 1.5. The correlation between RT-qPCR-based fold change and microarray-based fold change of *ABCA13* was  $r = 0.782$ ,  $p < 0.0001$  (Figure 4).

Due to the low number of significant DEGs (<10), functional enrichment and pathway enrichment could not be completed.

## DISCUSSION

This longitudinal study of HBOT in healthy aging subjects demonstrates that HBOT strongly associated with blood cells gene expression profiles, as estimated by RNA microarrays. To the best of our knowledge, this serves as the first study assessing human *in vivo* HBOT-mediated changes in RNA expression. Most of the significant changes occurred during the HBOT program and normalized two weeks post-HBOT. However, 19 genes were significantly different even two weeks after the last HBOT session.

Multiple previous studies have demonstrated the effects of HBOT on single gene products in isolated cells [17–25]. Recently, gene array analyses have demonstrated differential gene expression of isolated cells exposed to




Figure 1. Patient flowchart.

**Table 1. Baseline characteristics.**

<b>N</b>		27
<b>Age (years)</b>		70.39 ± 3.72
<b>BMI</b>		26.55 ± 3.06
<b>Males</b>		14 (51.9%)
<b>Complete blood count</b>		
	Hemoglobin	13.98 ± 1.39
	White blood cells	6.48 ± 1.22
	Platelets	245.05 ± 45.4
<b>Chronic medical conditions</b>		
	Atrial fibrillation	4 (14.8%)
	Hypothyroidism	4 (14.8%)
	Obstructive sleep apnea	3 (11.1%)
	Asthma	1 (3.7%)
	BPH	7 (25.9%)
	GERD	3 (11.1%)
	Osteoporosis	5 (18.5%)
	Rheumatic arthritis	0 (0%)
	Osteoarthritis	4 (14.8%)
	Diabetes mellitus	3 (11.1%)
	Hypertension	6 (22.2%)
	Dyslipidemia	14 (51.9%)
	Ischemic heart disease	0 (0%)
	History of smoking	9 (33.3%)
<b>Chronic medications</b>		
	Anti-aggregation	6 (22.2%)
	ACE-Inhibitors/ARB blockers	4 (14.8%)
	Beta blockers	4 (14.8%)
	Calcium blockers	2 (7.4%)
	Alpha blockers	5 (18.5%)
	Diuretics	2 (7.4%)
	Statins	8 (29.6%)
	Oral hypoglycemic	1 (3.7%)
	Bisphosphonates	1 (3.7%)
	Proton pump inhibitors	2 (7.4%)
	Hormones	2 (7.4%)
	Benzodiazepines	2 (7.4%)
	SSRI	5 (18.5%)

Abbreviations: BMI: body mass index; BPH: benign prostate hyperplasia; GERD: gastroesophageal reflux disease; ACE: angiotensin converting enzyme; ARB: angiotensin receptor blocker; SSRI: selective serotonin reuptake inhibitor.

either single hyperoxic exposure and/or high atmospheric pressure exposure [11–14]. Upregulated


genes included anti-inflammatory and growth and repair hormones genes, while downregulated genes included

**Table 2. Top DEG following the last HBOT session (fold change>1.5).**

ID	Baseline Avg (log2)	60th HBOT Avg (log2)	Fold Change	P-val	FDR Q-val	Gene Symbol	Description
TC0700007480.hg.1	8.39	7.2	-2.28	1.73E-13	1.86E-09	ABCA13	ATP Binding Cassette Subfamily A Member 13
TC0100008620.hg.1	11.85	10.74	-2.16	4.4E-12	3.15E-08	DNAJC6	DnaJ (Hsp40) homolog, subfamily C, member 6
TC0500009411.hg.1	8.67	7.86	-1.75	1.57E-10	2.60E-07	RANBP17	RAN binding protein 17
TC1100013133.hg.1	15.56	14.92	-1.56	7/62E-14	1.63E-09	HBG2; HBG1	hemoglobin, gamma G; hemoglobin, gamma A
TC1600007007.hg.1	12.09	11.48	-1.53	4.55E-11	1.08E-07	PDXDC1	pyridoxal-dependent decarboxylase domain containing 1

apoptosis and pro-inflammation genes. However, isolated cell studies have significant disadvantages. First, *in vitro* oxygen diffusion is considerably different than *in vivo* diffusion. Obtaining rapid changes in PO<sub>2</sub> within the cells/tissue culture medium proves to be difficult and limits a precise correlation between cellular processes/events and in environmental oxygen

concentration changes [26, 27]. Second, the oxygen partial pressure that cells are truly exposed *in vitro* is significantly different than what is used in *in vivo* studies [28, 29]. Third, *in vitro* studies lack the microsystem interactions which occur *in vivo*. Fourth, gene expression was evaluated following a single hyperoxic exposure. The current clinical study, provides



**Figure 2. Alterations in gene expression profile following the last HBOT session.** Volcano plot showing the distribution of gene expression following 60 HBOT sessions compared to baseline. Significance versus log<sub>2</sub> fold change is plotted on the y and x, respectively. Red dots represent the significant upregulated DEGs, green dots represent the significant downregulated DEGs.

**Table 3. Overlapping differentially expressed genes following the last session and two weeks post-HBOT.**


ID	Baseline Avg (log2)	60th HBOT Avg (log2)	2 weeks post HBOT Avg (log2)	60th session Fold Change	2 weeks post HBOT Fold Change	P-val	Q-val (FDR)	Gene Symbol	Description
TC0700007480.hg.1	8.39	7.2	7.76	-2.28	-1.54	1.73e-13	1.86E-09	ABCA13	ATP Binding Cassette Subfamily A Member 13

DEG, for the first time in humans, following 60 repeated hyperoxic exposures, both within 24 hours and two weeks after the last exposure.

HBOT is a well-established treatment modality for non-healing wounds, radiation injuries as well as different hypoxic or ischemic events (such as carbon monoxide toxicity, infections, etc.). In recent years, a growing body of evidence from preclinical as well as clinical trials demonstrates HBOT’s efficacy for neurological indications including post-stroke and post-

traumatic brain injury [30–36], idiopathic sudden sensorineural hearing loss [37] central sensitization syndromes such as fibromyalgia [38, 39], age related cognitive decline [10] and animal models of Alzheimer’s disease [40]. For the first time, the current study aimed to evaluate the effects of HBOT on the transcriptome of aging humans without any functional limiting diseases.

With regards to ageing, Peters et al. identified 1497 genes that are differentially expressed as we



**Figure 3. Alterations in gene expression profile two weeks after completion of 60 HBOT sessions.** Volcano plot showing the distribution of gene expression two weeks following 60 HBOT sessions compared to baseline. Significance versus log2 fold change is plotted on the y and x axes, respectively. Red dots represent the significant upregulated DEGs, green dots represent the significant downregulated DEGs.

chronologically age [2], out of which, 240 (16%) were differentially expressed in our study following the last HBOT session. For instance, polymorphism in the FOXO3 gene, which encodes forkhead box O-3 transcription factor has been associated with longevity in different human populations [41–43]. In the current study, FOXO3 gene expression changed by 1.22. RUNX3, a hematopoietic stem and progenitor cell factor whose levels decline with aging [44] increased by 1.29. It is important to note that these age genetic signatures generated by different groups show relatively little overlap with each other.

Whole blood transcriptomes are dynamic per definition and represent the cellular state at a certain point. Therefore, it is expected that most of the DEGs occurred during the repeated intermittent hyperbaric exposures, and gradually returned to their previous state two weeks following the last exposure. Previous studies have shown long-term systemic and cellular effects of HBOT, including angiogenesis, stem cells proliferation and mobilization and mitochondrial biogenesis [45] which may be partially explained by these DEGs. The effects last in the protein, tissue and system level

compared with the temporal transcription level. ABCA13 is a member of the ATP binding cassette gene subfamily A (ABCA). High expression of ABCA13 has been demonstrated in prostate cancer, leukemia colorectal cancer, as well as several tumor cell lines in central nervous system [46–48]. Additionally, both schizophrenia and bipolar disorder were associated with ABCA13 high expression [48]. Interestingly, expression increases following an ischemic stroke. Lastly, ABCA13 has been shown to decrease significantly in bone-marrow derived mesenchymal stem cells [49]. It has been previously shown HBOT can induce significant recruitment and migration of bone-marrow derived stem cells, both hematopoietic [50] and mesenchymal types [51], which may partially explain the significant down regulation of ABCA13. ABCA13 was the single DEG which remained significant even two weeks following the last HBOT sessions.

The main limitations of the study are related to the lack of a placebo group, required for definitive causality of the observed transcriptome changes, and the relatively small sample size. Additionally, further study with longer observation time would shed better light on




Figure 4. ABCA13 gene expression validation with PCR.



additional factors such as subject-specific expression levels differences and their relation to treatment response. These limitations are partially mitigated by the longitudinal design of the study with both the first and second baseline samples from each individual being combined to detect major changes associated with HBOT, rather than temporal/retest effects.

## CONCLUSIONS

For the first time in humans, the study provides direct evidence of HBOT is associated with transcriptome changes in whole blood samples. Our results have demonstrated significant changes in specific gene expressions of normal aging adults.

## AUTHOR CONTRIBUTIONS

All authors contributed substantially to the preparation of this manuscript. HA, FR and ES were responsible for protocol design. HA, ZY, BY, DKM and ES were responsible for patients' recruitment. HA, ZY, BY, DKM, FR and VD were responsible for data acquisition. HA, FR, VD, CM and ES were responsible for data analysis. All authors revised and finalized the manuscript.

## ACKNOWLEDGMENTS

We would like to thank Dr. Mechael Kanovsky for his editing of this manuscript.

## CONFLICTS OF INTEREST

AH, BY, ZY work for AVIV Scientific LTD. ES is a shareholder at AVIV Scientific LTD.

## FUNDING

The study was funded by a research grant from the Sagol network for neuroscience established by Mr. Sami Sagol.

## REFERENCES

1. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell*. 2013; 153:1194–217.  
<https://doi.org/10.1016/j.cell.2013.05.039>  
PMID:[23746838](https://pubmed.ncbi.nlm.nih.gov/23746838/)
2. Peters MJ, Joehanes R, Pilling LC, Schurmann C, Conneely KN, Powell J, Reinmaa E, Sutphin GL, Zhernakova A, Schramm K, Wilson YA, Kobes S, Tukiainen T, et al, and NABEC/UKBEC Consortium. The transcriptional landscape of age in human peripheral blood. *Nat Commun*. 2015; 6:8570.  
<https://doi.org/10.1038/ncomms9570>  
PMID:[26490707](https://pubmed.ncbi.nlm.nih.gov/26490707/)
3. Barth E, Srivastava A, Stojiljkovic M, Frahm C, Axer H, Witte OW, Marz M. Conserved aging-related signatures of senescence and inflammation in different tissues and species. *Aging (Albany NY)*. 2019; 11:8556–72.  
<https://doi.org/10.18632/aging.102345>  
PMID:[31606727](https://pubmed.ncbi.nlm.nih.gov/31606727/)
4. Calabria E, Mazza EM, Dyar KA, Pogliaghi S, Bruseghini P, Morandi C, Salvagno GL, Gelati M, Guidi GC, Bicciato S, Schiaffino S, Schena F, Capelli C. Aging: a portrait from gene expression profile in blood cells. *Aging (Albany NY)*. 2016; 8:1802–21.  
<https://doi.org/10.18632/aging.101016>  
PMID:[27545843](https://pubmed.ncbi.nlm.nih.gov/27545843/)
5. Cimino F, Balestra C, Germonpré P, De Bels D, Tillmans F, Saija A, Speciale A, Virgili F. Pulsed high oxygen induces a hypoxic-like response in human umbilical endothelial cells and in humans. *J Appl Physiol (1985)*. 2012; 113:1684–89.  
<https://doi.org/10.1152/jappphysiol.00922.2012>  
PMID:[23042909](https://pubmed.ncbi.nlm.nih.gov/23042909/)
6. Sunkari VG, Lind F, Botusan IR, Kashif A, Liu ZJ, Ylä-Herttuala S, Brismar K, Velazquez O, Catrina SB. Hyperbaric oxygen therapy activates hypoxia-inducible factor 1 (HIF-1), which contributes to improved wound healing in diabetic mice. *Wound Repair Regen*. 2015; 23:98–103.  
<https://doi.org/10.1111/wrr.12253>  
PMID:[25532619](https://pubmed.ncbi.nlm.nih.gov/25532619/)
7. Milovanova TN, Bhopale VM, Sorokina EM, Moore JS, Hunt TK, Hauer-Jensen M, Velazquez OC, Thom SR. Hyperbaric oxygen stimulates vasculogenic stem cell growth and differentiation in vivo. *J Appl Physiol (1985)*. 2009; 106:711–28.  
<https://doi.org/10.1152/jappphysiol.91054.2008>  
PMID:[19023021](https://pubmed.ncbi.nlm.nih.gov/19023021/)
8. Yang Y, Wei H, Zhou X, Zhang F, Wang C. Hyperbaric oxygen promotes neural stem cell proliferation by activating vascular endothelial growth factor/extracellular signal-regulated kinase signaling after traumatic brain injury. *Neuroreport*. 2017; 28:1232–38.  
<https://doi.org/10.1097/WNR.0000000000000901>  
PMID:[28953090](https://pubmed.ncbi.nlm.nih.gov/28953090/)
9. Fife CE, Eckert KA, Carter MJ. An Update on the Appropriate Role for Hyperbaric Oxygen: Indications and Evidence. *Plast Reconstr Surg*. 2016; 138:107S–16S.  
<https://doi.org/10.1097/PRS.0000000000002714>  
PMID:[27556750](https://pubmed.ncbi.nlm.nih.gov/27556750/)

10. Hadanny A, Daniel-Kotovskiy M, Suzin G, Boussi-Gross R, Catalogna M, Dagan K, Hachmo Y, Abu Hamed R, Sasson E, Fishlev G, Lang E, Polak N, Doeniyas K, et al. Cognitive enhancement of healthy older adults using hyperbaric oxygen: a randomized controlled trial. *Aging (Albany NY)*. 2020; 12:13740–61.  
<https://doi.org/10.18632/aging.103571>  
PMID:[32589613](https://pubmed.ncbi.nlm.nih.gov/32589613/)
11. Oh S, Lee E, Lee J, Lim Y, Kim J, Woo S. Comparison of the effects of 40% oxygen and two atmospheric absolute air pressure conditions on stress-induced premature senescence of normal human diploid fibroblasts. *Cell Stress Chaperones*. 2008; 13:447–58.  
<https://doi.org/10.1007/s12192-008-0041-5>  
PMID:[18465208](https://pubmed.ncbi.nlm.nih.gov/18465208/)
12. Chen Y, Nadi NS, Chavko M, Auker CR, McCarron RM. Microarray analysis of gene expression in rat cortical neurons exposed to hyperbaric air and oxygen. *Neurochem Res*. 2009; 34:1047–56.  
<https://doi.org/10.1007/s11064-008-9873-8>  
PMID:[19015983](https://pubmed.ncbi.nlm.nih.gov/19015983/)
13. Godman CA, Chheda KP, Hightower LE, Perdrizet G, Shin DG, Giardina C. Hyperbaric oxygen induces a cytoprotective and angiogenic response in human microvascular endothelial cells. *Cell Stress Chaperones*. 2010; 15:431–42.  
<https://doi.org/10.1007/s12192-009-0159-0>  
PMID:[19949909](https://pubmed.ncbi.nlm.nih.gov/19949909/)
14. Kendall AC, Whatmore JL, Harries LW, Winyard PG, Eggleton P, Smerdon GR. Different oxygen treatment pressures alter inflammatory gene expression in human endothelial cells. *Undersea Hyperb Med*. 2013; 40:115–23.  
PMID:[23682543](https://pubmed.ncbi.nlm.nih.gov/23682543/)
15. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B (Methodological)*. 1995; 57:289–300.
16. Shamir R, Maron-Katz A, Tanay A, Linhart C, Steinfeld I, Sharan R, Shiloh Y, Elkon R. EXPANDER—an integrative program suite for microarray data analysis. *BMC Bioinformatics*. 2005; 6:232.  
<https://doi.org/10.1186/1471-2105-6-232>  
PMID:[16176576](https://pubmed.ncbi.nlm.nih.gov/16176576/)
17. Ishii Y, Miyanaga Y, Shimojo H, Ushida T, Tateishi T. Effects of hyperbaric oxygen on procollagen messenger RNA levels and collagen synthesis in the healing of rat tendon laceration. *Tissue Eng*. 1999; 5:279–86.  
<https://doi.org/10.1089/ten.1999.5.279>  
PMID:[10434074](https://pubmed.ncbi.nlm.nih.gov/10434074/)
18. Yang JT, Chang CN, Lee TH, Lin TN, Hsu JC, Hsu YH, Wu JH. Hyperbaric oxygen treatment decreases post-ischemic neurotrophin-3 mRNA down-regulation in the rat hippocampus. *Neuroreport*. 2001; 12:3589–92.  
<https://doi.org/10.1097/00001756-200111160-00043>  
PMID:[11733717](https://pubmed.ncbi.nlm.nih.gov/11733717/)
19. Lin S, Shyu KG, Lee CC, Wang BW, Chang CC, Liu YC, Huang FY, Chang H. Hyperbaric oxygen selectively induces angiopoietin-2 in human umbilical vein endothelial cells. *Biochem Biophys Res Commun*. 2002; 296:710–15.  
[https://doi.org/10.1016/s0006-291x\(02\)00924-5](https://doi.org/10.1016/s0006-291x(02)00924-5)  
PMID:[12176040](https://pubmed.ncbi.nlm.nih.gov/12176040/)
20. Freiberger J, Coulombe K, Suliman H, Carraway M, Piantadosi C. Superoxide dismutase responds to hyperoxia in rat hippocampus. *Undersea Hyperb Med*. 2004; 31:227–32.  
PMID:[15485085](https://pubmed.ncbi.nlm.nih.gov/15485085/)
21. Yu Y, Matsuyama Y, Yanase M, Ito S, Adachi K, Satake K, Ishiguro N, Kiuchi K. Effects of hyperbaric oxygen on GDNF expression and apoptosis in spinal cord injury. *Neuroreport*. 2004; 15:2369–73.  
<https://doi.org/10.1097/00001756-200410250-00014>  
PMID:[15640758](https://pubmed.ncbi.nlm.nih.gov/15640758/)
22. Takeyama N, Sakai H, Ohtake H, Mashitori H, Tamai K, Saotome K. Effects of hyperbaric oxygen on gene expressions of procollagen, matrix metalloproteinase and tissue inhibitor of metalloproteinase in injured medial collateral ligament and anterior cruciate ligament. *Knee Surg Sports Traumatol Arthrosc*. 2007; 15:443–52.  
<https://doi.org/10.1007/s00167-006-0241-4>  
PMID:[17187281](https://pubmed.ncbi.nlm.nih.gov/17187281/)
23. Sun L, Marti HH, Veltkamp R. Hyperbaric oxygen reduces tissue hypoxia and hypoxia-inducible factor-1 alpha expression in focal cerebral ischemia. *Stroke*. 2008; 39:1000–1006.  
<https://doi.org/10.1161/STROKEAHA.107.490599>  
PMID:[18239183](https://pubmed.ncbi.nlm.nih.gov/18239183/)
24. Romero-Valdovinos M, Cárdenas-Mejía A, Gutiérrez-Gómez C, Flisser A, Kawa-Karasik S, Ortiz-Monasterio F. Keloid skin scars: the influence of hyperbaric oxygenation on fibroblast growth and on the expression of messenger RNA for insulin like growth factor and for transforming growth factor. *In Vitro Cell Dev Biol Anim*. 2011; 47:421–24.  
<https://doi.org/10.1007/s11626-011-9418-3>  
PMID:[21573718](https://pubmed.ncbi.nlm.nih.gov/21573718/)
25. Gajendrareddy PK, Sen CK, Horan MP, Marucha PT. Hyperbaric oxygen therapy ameliorates stress-impaired dermal wound healing. *Brain Behav Immun*. 2005; 19:217–22.

- <https://doi.org/10.1016/j.bbi.2004.09.003>  
PMID:[15797310](https://pubmed.ncbi.nlm.nih.gov/15797310/)
26. Allen CB, Schneider BK, White CW. Limitations to oxygen diffusion and equilibration in in vitro cell exposure systems in hyperoxia and hypoxia. *Am J Physiol Lung Cell Mol Physiol*. 2001; 281:L1021–27.  
<https://doi.org/10.1152/ajplung.2001.281.4.L1021>  
PMID:[11557606](https://pubmed.ncbi.nlm.nih.gov/11557606/)
27. Place TL, Domann FE, Case AJ. Limitations of oxygen delivery to cells in culture: An underappreciated problem in basic and translational research. *Free Radic Biol Med*. 2017; 113:311–22.  
<https://doi.org/10.1016/j.freeradbiomed.2017.10.003>  
PMID:[29032224](https://pubmed.ncbi.nlm.nih.gov/29032224/)
28. Ortiz-Prado E, Dunn JF, Vasconez J, Castillo D, Viscor G. Partial pressure of oxygen in the human body: a general review. *Am J Blood Res*. 2019; 9:1–14.  
PMID:[30899601](https://pubmed.ncbi.nlm.nih.gov/30899601/)
29. Carreau A, El Hafny-Rahbi B, Matejuk A, Grillon C, Kieda C. Why is the partial oxygen pressure of human tissues a crucial parameter? Small molecules and hypoxia. *J Cell Mol Med*. 2011; 15:1239–53.  
<https://doi.org/10.1111/j.1582-4934.2011.01258.x>  
PMID:[21251211](https://pubmed.ncbi.nlm.nih.gov/21251211/)
30. Boussi-Gross R, Golan H, Fishlev G, Bechor Y, Volkov O, Bergan J, Friedman M, Hoofien D, Shlamkovitch N, Ben-Jacob E, Efrati S. Hyperbaric oxygen therapy can improve post concussion syndrome years after mild traumatic brain injury - randomized prospective trial. *PLoS One*. 2013; 8:e79995.  
<https://doi.org/10.1371/journal.pone.0079995>  
PMID:[24260334](https://pubmed.ncbi.nlm.nih.gov/24260334/)
31. Efrati S, Fishlev G, Bechor Y, Volkov O, Bergan J, Kliakhandler K, Kamiager I, Gal N, Friedman M, Ben-Jacob E, Golan H. Hyperbaric oxygen induces late neuroplasticity in post stroke patients--randomized, prospective trial. *PLoS One*. 2013; 8:e53716.  
<https://doi.org/10.1371/journal.pone.0053716>  
PMID:[23335971](https://pubmed.ncbi.nlm.nih.gov/23335971/)
32. Mukherjee A, Raison M, Sahni T, Arya A, Lambert J, Marois P, James PB, Parent A, Ballaz L. Intensive rehabilitation combined with HBO2 therapy in children with cerebral palsy: a controlled longitudinal study. *Undersea Hyperb Med*. 2014; 41:77–85.  
PMID:[24851544](https://pubmed.ncbi.nlm.nih.gov/24851544/)
33. Hadanny A, Golan H, Fishlev G, Bechor Y, Volkov O, Suzin G, Ben-Jacob E, Efrati S. Hyperbaric oxygen can induce neuroplasticity and improve cognitive functions of patients suffering from anoxic brain damage. *Restor Neurol Neurosci*. 2015; 33:471–86.  
<https://doi.org/10.3233/RNN-150517>  
PMID:[26409406](https://pubmed.ncbi.nlm.nih.gov/26409406/)
34. Tal S, Hadanny A, Berkovitz N, Sasson E, Ben-Jacob E, Efrati S. Hyperbaric oxygen may induce angiogenesis in patients suffering from prolonged post-concussion syndrome due to traumatic brain injury. *Restor Neurol Neurosci*. 2015; 33:943–51.  
<https://doi.org/10.3233/RNN-150585>  
PMID:[26484702](https://pubmed.ncbi.nlm.nih.gov/26484702/)
35. Hadanny A, Rittblat M, Bitterman M, May-Raz I, Suzin G, Boussi-Gross R, Zemel Y, Bechor Y, Catalogna M, Efrati S. Hyperbaric oxygen therapy improves neurocognitive functions of post-stroke patients - a retrospective analysis. *Restor Neurol Neurosci*. 2020; 38:93–107.  
<https://doi.org/10.3233/RNN-190959>  
PMID:[31985478](https://pubmed.ncbi.nlm.nih.gov/31985478/)
36. Tal S, Hadanny A, Sasson E, Suzin G, Efrati S. Hyperbaric Oxygen Therapy Can Induce Angiogenesis and Regeneration of Nerve Fibers in Traumatic Brain Injury Patients. *Front Hum Neurosci*. 2017; 11:508.  
<https://doi.org/10.3389/fnhum.2017.00508>  
PMID:[29097988](https://pubmed.ncbi.nlm.nih.gov/29097988/)
37. Le W. *Hyperbaric Oxygen Therapy Indications*. UHMS. 2008; 12th edition:215–18.
38. Efrati S, Golan H, Bechor Y, Faran Y, Daphna-Tekoah S, Sekler G, Fishlev G, Ablin JN, Bergan J, Volkov O, Friedman M, Ben-Jacob E, Buskila D. Hyperbaric oxygen therapy can diminish fibromyalgia syndrome--prospective clinical trial. *PLoS One*. 2015; 10:e0127012.  
<https://doi.org/10.1371/journal.pone.0127012>  
PMID:[26010952](https://pubmed.ncbi.nlm.nih.gov/26010952/)
39. Hadanny A, Bechor Y, Catalogna M, Daphna-Tekoah S, Sigal T, Cohenpour M, Lev-Wiesel R, Efrati S. Hyperbaric Oxygen Therapy Can Induce Neuroplasticity and Significant Clinical Improvement in Patients Suffering From Fibromyalgia With a History of Childhood Sexual Abuse--Randomized Controlled Trial. *Front Psychol*. 2018; 9:2495.  
<https://doi.org/10.3389/fpsyg.2018.02495>  
PMID:[30618929](https://pubmed.ncbi.nlm.nih.gov/30618929/)
40. Shapira R, Efrati S, Ashery U. Hyperbaric oxygen therapy as a new treatment approach for Alzheimer's disease. *Neural Regen Res*. 2018; 13:817–18.  
<https://doi.org/10.4103/1673-5374.232475>  
PMID:[29863011](https://pubmed.ncbi.nlm.nih.gov/29863011/)
41. Flachsbarth F, Dose J, Gentschew L, Geismann C, Caliebe A, Knecht C, Nygaard M, Badarinarayan N, ElSharawy A, May S, Luzius A, Torres GG, Jentzsch M, et al. Identification and characterization of two functional variants in the human longevity gene FOXO3. *Nat Commun*. 2017; 8:2063.  
<https://doi.org/10.1038/s41467-017-02183-y>  
PMID:[29234056](https://pubmed.ncbi.nlm.nih.gov/29234056/)

42. Flachsbar F, Caliebe A, Kleindorp R, Blanché H, von Eller-Eberstein H, Nikolaus S, Schreiber S, Nebel A. Association of FOXO3A variation with human longevity confirmed in German centenarians. *Proc Natl Acad Sci U S A*. 2009; 106:2700–705. <https://doi.org/10.1073/pnas.0809594106> PMID:19196970
43. Soerensen M, Dato S, Christensen K, McGue M, Stevnsner T, Bohr VA, Christiansen L. Replication of an association of variation in the FOXO3A gene with human longevity using both case-control and longitudinal data. *Aging Cell*. 2010; 9:1010–17. <https://doi.org/10.1111/j.1474-9726.2010.00627.x> PMID:20849522
44. Balogh P, Adelman ER, Pluvinage JV, Capaldo BJ, Freeman KC, Singh S, Elagib KE, Nakamura Y, Kurita R, Sashida G, Zunder ER, Li H, Gru AA, et al. *RUNX3* levels in human hematopoietic progenitors are regulated by aging and dictate erythroid-myeloid balance. *Haematologica*. 2020; 105:905–13. <https://doi.org/10.3324/haematol.2018.208918> PMID:31171641
45. Hadanny A, Efrati S. The Hyperoxic-Hypoxic Paradox. *Biomolecules*. 2020; 10:958. <https://doi.org/10.3390/biom10060958> PMID:32630465
46. Albrecht C, Viturro E. The ABCA subfamily--gene and protein structures, functions and associated hereditary diseases. *Pflugers Arch*. 2007; 453:581–89. <https://doi.org/10.1007/s00424-006-0047-8> PMID:16586097
47. Hlavata I, Mohelnikova-Duchonova B, Vaclavikova R, Liska V, Pitule P, Novak P, Bruha J, Vycital O, Holubec L, Treska V, Vodicka P, Soucek P. The role of ABC transporters in progression and clinical outcome of colorectal cancer. *Mutagenesis*. 2012; 27:187–96. <https://doi.org/10.1093/mutage/ger075> PMID:22294766
48. Knight HM, Pickard BS, Maclean A, Malloy MP, Soares DC, McRae AF, Condie A, White A, Hawkins W, McGhee K, van Beck M, MacIntyre DJ, Starr JM, et al. A cytogenetic abnormality and rare coding variants identify ABCA13 as a candidate gene in schizophrenia, bipolar disorder, and depression. *Am J Hum Genet*. 2009; 85:833–46. <https://doi.org/10.1016/j.ajhg.2009.11.003> PMID:19944402
49. Barbet R, Peiffer I, Hutchins JR, Hatzfeld A, Garrido E, Hatzfeld JA. Expression of the 49 human ATP binding cassette (ABC) genes in pluripotent embryonic stem cells and in early- and late-stage multipotent mesenchymal stem cells: possible role of ABC plasma membrane transporters in maintaining human stem cell pluripotency. *Cell Cycle*. 2012; 11:1611–20. <https://doi.org/10.4161/cc.20023> PMID:22456339
50. Thom SR, Bhopale VM, Velazquez OC, Goldstein LJ, Thom LH, Buerk DG. Stem cell mobilization by hyperbaric oxygen. *Am J Physiol Heart Circ Physiol*. 2006; 290:H1378–86. <https://doi.org/10.1152/ajpheart.00888.2005> PMID:16299259
51. Peña-Villalobos I, Casanova-Maldonado I, Lois P, Prieto C, Pizarro C, Lattus J, Osorio G, Palma V. Hyperbaric Oxygen Increases Stem Cell Proliferation, Angiogenesis and Wound-Healing Ability of WJ-MSCs in Diabetic Mice. *Front Physiol*. 2018; 9:995. <https://doi.org/10.3389/fphys.2018.00995> PMID:30104981

## SUPPLEMENTARY MATERIALS

### Supplementary Tables

Please browse Full Text version to see the data of Supplementary Table 1.

**Supplementary Table 1. List of differentially expressed genes following the last HBOT session.**

**Supplementary Table 2. List of differentially expressed genes two weeks following the last HBOT session.**

<b>ID</b>	<b>Baseline Avg (log2)</b>	<b>2 weeks post HBOT Avg (log2)</b>	<b>Fold Change</b>	<b>FDR P-val</b>
TC0700007480.hg.1	8.39	7.76	-1.54	0.0079
TC0600007092.hg.1	12.17	11.87	-1.23	0.0029
TC0300009147.hg.1	14.86	14.62	-1.18	0.0004
TC0300008355.hg.1	4.66	4.47	-1.14	0.0066
TC0300010643.hg.1	11.54	11.36	-1.14	0.008
TC1400010619.hg.1	14.01	13.84	-1.13	0.0027
TC0200013096.hg.1	13.78	13.67	-1.08	0.0024
TC0200014971.hg.1	13.56	13.47	-1.07	0.009
TC1300008837.hg.1	12.61	12.76	1.11	0.008
TC1100013037.hg.1	9.74	9.9	1.12	0.006
TC0900009164.hg.1	9.99	10.17	1.13	0.0058
TC1500007093.hg.1	10.92	11.1	1.14	0.009
TC0800011171.hg.1	12.39	12.59	1.15	0.0027
TC0700013591.hg.1	12.34	12.55	1.16	0.0051
TC0800012147.hg.1	11.28	11.51	1.17	0.0008
TC2000008230.hg.1	8.56	8.79	1.17	0.0058
TC0300007387.hg.1	9.39	9.66	1.2	0.0058
TC0600014111.hg.1	8.26	8.54	1.21	0.0004
TC0800007004.hg.1	7.19	7.48	1.22	0.0051