

SUPPLEMENTARY TABLE

Supplementary Table 1. Sensitivity analysis: association of baseline RDW levels with recurrent ischemic stroke.

	Crude model		Adjusted model*	
	HR (95%CI)	P value	HR (95%CI)	P value
Model A				
Quartile				
Q1 (<12.4)	Ref		Ref	
Q2 (12.4 to 12.8)	1.003 (0.937-1.073)	0.932	1.003 (0.936-1.075)	0.927
Q3 (12.8 to 13.3)	1.169 (1.069-1.285)	0.076	1.166 (1.066-1.286)	0.050
Q4 (>13.3)	1.222 (1.091-1.369)	0.001	1.211 (1.075-1.364)	0.002
P for trend		<0.001		<0.001
Categories				
Q1-Q2 (<12.8)	Ref		Ref	
Q3-Q4 (>12.8)	1.305 (1.142-1.491)	<0.001	1.313 (1.144-1.506)	<0.001
Model B				
Quartile				
Q1 (<12.4)	Ref		Ref	
Q2 (12.4 to 12.8)	1.003 (0.938-1.073)	0.921	1.003 (0.937-1.074)	0.932
Q3 (12.8 to 13.3)	1.225 (1.006-1.505)	0.044	1.248 (1.011-1.541)	0.039
Q4 (>13.3)	1.289 (1.130-1.470)	<0.001	1.293 (1.127-1.485)	<0.001
P for trend		<0.001		<0.001
Categories				
Q1-Q2 (<12.8)	Ref		Ref	
Q3-Q4 (>12.8)	1.548 (1.319-1.818)	<0.001	1.581 (1.340-1.864)	<0.001

*Adjusted for age, sex, hypertension, diabetes mellitus, smoking, drinking, body mass index, and coronary heart disease. Q, quartile. Model A: all the lost to follow up patients (n=175) were considered as stroke recurrence. Model B: all the lost to follow up patients (n=175) were considered as non-stroke recurrence.