

Correction

Correction for: Posttranscriptional regulation of AKT by circular RNA angiomin-1 mediates chemoresistance against paclitaxel in breast cancer cells

Jian Ma^{1,2}, Ling Fang¹, Qi Yang^{1,3}, Steven Hibberd¹, William W. Du^{1,4}, Nan Wu¹, Burton B. Yang^{1,4}

¹Sunnybrook Research Institute, Sunnybrook Health Sciences Centre, Toronto, Canada

²Department of Urology, The Affiliated Yantai Yuhuangding Hospital of Qingdao University, Yantai, China

³The First Hospital, Jilin University, Jilin, China

⁴Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Canada

Correspondence to: Burton B. Yang, Nan Wu; email: byang@sri.utoronto.ca, wunan.my@gmail.com

Keywords: circular RNA, circAMOTL1, chemoresistance, PAX, AKT

Abbreviations: DMEM: Dulbecco's modified Eagle's medium; FBS: fetal bovine serum; GFP: green fluorescent protein

Original article: *Aging (Albany NY)* 2019; 11:11369–11381

PMID: [31819016](https://pubmed.ncbi.nlm.nih.gov/31819016/)

PMCID: [PMC6932896](https://pubmed.ncbi.nlm.nih.gov/PMC6932896/)

doi: [10.18632/aging.102535](https://doi.org/10.18632/aging.102535)

This article has been corrected: The authors have requested a replacement for Figure 4D. A representative image of the si-circAMOTL1 Control sample in Figure 4D was mistakenly mixed up. The correct panel has now been prepared with the si-circAMOTL1 Control image from the original set of experiments. This alteration does not affect the results or conclusions of this work.

The correct Figure 4D is presented below:




Figure 4. Effect of circAMOTL1 siRNA on PAX treatment. MDA-MB-231 cells transfected with negative control or circAMOTL1 siRNA (si-circAMOTL1) were treated with 1 µg/ml PAX for 24 hours. ... ((D) Cell invasive ability upon PAX treatment was measured with Matrigel invasion assay. Experiments were performed in triplicate. ** $p < 0.01$ compared to untreated negative control. ## $p < 0.01$ compared to untreated si-circAMOTL1.