

## SUPPLEMENTARY TABLE

**Supplementary Table 2. Differential expression status of m6A regulatory genes in TCGA\_PAAD patients compared with GTEx normal individuals.**

|         | log2FoldChange* | p-value |
|---------|-----------------|---------|
| YTHDC2  | -0.099          | 0.012   |
| METTL14 | -0.120          | 0.024   |
| YTHDC1  | -0.113          | 0.045   |
| IGF2BP2 | 0.341           | 0.048   |
| ZCCHC4  | 0.103           | 0.078   |
| IGF2BP3 | 0.166           | 0.389   |
| FTO     | 0.038           | 0.394   |
| YTHDF2  | -0.090          | 0.403   |
| YTHDF3  | -0.031          | 0.479   |
| ZC3H13  | 0.063           | 0.483   |
| RBM15   | 0.064           | 0.574   |
| YTHDF1  | -0.062          | 0.672   |
| WTAP    | -0.036          | 0.774   |
| METTL3  | -0.027          | 0.816   |
| ALKBH5  | -0.012          | 0.861   |
| IGF2BP1 | 0.040           | 0.876   |
| YTHDC2  | -0.099          | 0.012   |
| METTL14 | -0.120          | 0.024   |

\*log2FoldChange < 0 meant that a certain gene was down-expressed in PAAD tissue than that in normal.