

SUPPLEMENTARY TABLE

Supplementary Table 1. Clinical features of 36 patients with HCC.

Variable	Number/Mean±SD
Age (year)	52.61 ± 12.59
Gender (male/female)	32/4
Family history of HCC (+/-)	5/31
History of drinking (+/-)	6/30
Cirrhosis (+/-)	27/9
HBV DNA (\geq / $<$ 100 copies/ml)	15/21
HBsAg (+/-)	30/6
HBsAb (+/-)	4/32
HBcAb (+/-)	36/0
HBeAg (+/-)	2/34
HBeAb (+/-)	4/32
alpha-Fetoproteins ($>$ 20/ \leq 20 μ g/L)	18/18
Alanine aminotransferase ($>$ 40/ \leq 40 U/L)	15/21
Aspartate aminotransferase ($>$ 40/ \leq 40 U/L)	20/16
Albumin ($>$ 35/ \leq 35 g/L)	22/14
Total bilirubin ($>$ 17/ \leq 17 μ mol/L)	14/22
γ -glutamyl transpeptidase ($>$ 100/ \leq 100 U/L)	27/9
Tumor size ($>$ 3/ \leq 3 cm)	28/8
Number of tumor nodules ($>$ 1/ \leq 1)	7/29
Portal vein thrombus (+/-)	5/31
Distance metastasis (+/-)	0/36
Pathological differentiation (high/moderate/low)	1/15/20
TNM stage (I/II/III/IIIa/IIIb)	8/4/19/5

SD, standard deviation.