

SUPPLEMENTARY TABLE

Supplementary Table 1. Baseline comparison of patients with included and excluded.

Characteristics	Included patients	Excluded patients	P-value
Demography and clinical features			
Age(yr), mean \pm SD	61.88 \pm 11.11	62.76 \pm 11.57	<.0001
Male, <i>n</i> (%)	6343 (69.7)	4021 (66.4)	<.0001
Smoking, <i>n</i> (%)	3317 (36.4)	2095 (34.6)	0.0201
Drinking, <i>n</i> (%)	3553 (39.0)	2162 (35.7)	<.0001
Medical history			
Ischemic stroke, <i>n</i> (%)	1850 (20.3)	1299 (21.4)	0.0943
Coronary heart diseases, <i>n</i> (%)	914 (10.0)	694 (11.5)	0.0055
Atrial fibrillation, <i>n</i> (%)	592 (6.5)	427 (7.1)	0.1876
Hypertension, <i>n</i> (%)	5717 (62.8)	3777 (62.3)	0.5843
Diabetes mellitus, <i>n</i> (%)	2126 (23.3)	1384 (22.8)	0.4722
Hypercholesterolemia, <i>n</i> (%)	696 (7.6)	495 (8.2)	0.2372
Hours of event onset, median (IQR)	10 (3–25)	20 (3–76)	<.0001
NIHSS at admission, median (IQR)	3 (2–6)	3 (1–6)	<.0001
mRS score before onset \geq3, <i>n</i> (%)	370 (4.1)	287 (4.7)	0.0458
Laboratory tests			
WBC, ($10^9/L$)	7.24 \pm 2.21	7.91 \pm 40.32	0.0561
Neutrophil, ($10^9/L$)	4.85 \pm 2.23	4.99 \pm 2.32	0.0463
Platelet, ($10^9/L$)	217.60 \pm 63.37	217.32 \pm 65.61	0.2199
Lymphocyte, ($10^9/L$)	1.81 \pm 0.79	1.78 \pm 0.84	0.0106
FPG (mmol/L)	6.46 \pm 2.62	6.34 \pm 2.49	0.0210
LDL (mmol/L)	2.56 \pm 1.04	2.55 \pm 1.06	0.1947
HCY (μ mol/L)	18.93 \pm 12.26	19.24 \pm 12.47	0.0425
hs-CRP (mg/L)	5.67 \pm 18.22	10.74 \pm 33.76	<.0001

Abbreviations: SD: standard deviation; IQR: interquartile range; NIHSS: National Institutes of Health Stroke Scale; Q: quartile; mRS: modified Rankin Scale; WBC: white blood cell; FPG: fasting plasma glucose; LDL: low-density lipoprotein; Hcy: homocysteine; hs-CRP: high sensitivity C reactive protein.