

Supplementary Table 1. The qRT-PCR primers and probes.

Gene symbol	Forward primer	Probe	Reverse primer
<i>CROP</i>	5'-CAACGTCGACA ATTGAAAGC-3'	5'-FAM-ACCCGGGACTG GGCATCTCC-TAMRA-3'	5'-GCATAGCCCAT GTGTTGTTT-3'
<i>NFIB</i>	5'-CTTATGACCCAT CCAGTCCA-3'	5'-FAM-CCAGCCAGTCCTGGT ACCTGGG-TAMRA-3'	5'-TGACACTTGGA AAGGAACCA-3'
<i>NKTR</i>	5'-TGCAGAACCTG AACCGAAGAT-3'	5'-FAM-CCTGATGTTGCA CCCATTGTAAGTGATCA G-TAMRA-3'	5'-CGTCCAGACTT TGATACAGATGGT-3'
<i>MALAT1</i>	5'-GGAAGTAATTCA AGATCAAGA-3'	5'-FAM-CCTGTCTGCTGC TGTCTTCCTA-TAMRA-3'	5'-CAGTCCTAGCT TCATCAA-3'
<i>β-ACTIN</i>	5'-GGTCATCACCAT TGGCAATG-3'	5'-FAM-CGGTTCCGCTGCCCT GAGGC-TAMRA-3'	5'-TCCATGCCAG GAAGGAA-3'

FAM, 6-carboxyfluorescein; TAMRA, 6-carboxytetramethylrhodamine.